

ACTIMMUNE

MEDICATION(S)

ACTIMMUNE

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

1. Hypersensitivity to Actimmune or E. coli derived products

REQUIRED MEDICAL INFORMATION

1. Dx chronic granulomatous disease OR Dx severe malignant osteopetrosis

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

BvD

PART B PREREQUISITE

N/A

AKEEGA

MEDICATION(S)

AKEEGA

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of deleterious or suspected deleterious BRCA-mutated (BRCAm) metastatic castration-resistant prostate cancer (mCRPC) 2. Contraindication, intolerance, or failure of Lynparza

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

ALECENSA

MEDICATION(S)

ALECENSA

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of NSCLC that is ALK-positive
 - a. Tumors are anaplastic lymphoma kinase (ALK)-positive

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

ALS

MEDICATION(S)

RADICAVA ORS, RADICAVA ORS STARTER KIT

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1)Dx of amyotrophic lateral sclerosis (ALS) as defined by the revised El Escorial criteria

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

ALUNBRIG

MEDICATION(S)

ALUNBRIG

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

- 1)Dx of metastatic non-small cell lung cancer (NSCLC)
 - a)Tumors are anaplastic lymphoma kinase (ALK)-positive
- 2)Patient has failed/intolerance/contraindication to Alecensa

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

ANDROGENS

MEDICATION(S)

TESTOSTERONE 10 MG/ACT (2%) GEL, TESTOSTERONE 30 MG/ACT SOLUTION

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Two separate testosterone levels drawn on different dates where the serum testosterone level is below 300 ng/dL, 2. Pt experiences at least ONE of the following: a. malaise, b. fatigue, c. lethargy, d. muscle loss, e. depression, f. decreased libido,

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

ANTIPSYCHOTIC

MEDICATION(S)

CAPLYTA, COBENFY, COBENFY STARTER PACK, FANAPT, FANAPT TITRATION PACK, LYBALVI, VERSACLOZ, VRAYLAR 1.5 MG CAP, VRAYLAR 3 MG CAP, VRAYLAR 4.5 MG CAP, VRAYLAR 6 MG CAP

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Previous trial on at least ONE of the following: aripiprazole, clozapine, fluoxetine-olanzapine, haloperidol, olanzapine, quetiapine, risperidone, ziprasidone 2. For Major Depressive Disorder (MDD) or Schizophrenia: Previous trial on Rexulti

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

APO B

MEDICATION(S)

JUXTAPID

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

Pt has none of the following health conditions or health concerns: a. History of significant hepatic disease, b. Alcohol abuse,

REQUIRED MEDICAL INFORMATION

1. Pt has untreated, fasting LDL cholesterol greater than 500 mg/dL AND triglycerides less than 300 mg/dL, 2. Pt meets a OR b AND c of the following: a. Pt has documented mutations in both alleles of the LDL receptor or of other genes known to affect LDL receptor function, b. Both of pt's parents have a hx of untreated total cholesterol of greater than 250 mg/dL, c. Pt has xanthomas present before age 10, 3. Pt has failed or is currently taking at least ONE of the following: a. Atorvastatin, Rosuvastatin, or Simvastatin b. Has documented intolerance (e.g. rhabdomyolysis) to statin therapy 4. Pt has previous trial of Repatha OR Praluent

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

1. Initial: 6 mo, 2. Reauthorization: Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

AQNEURSA

MEDICATION(S)

AQNEURSA

PENDING CMS APPROVAL

ARCALYST

MEDICATION(S)

ARCALYST

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

1. Combination use with a TNF-inhibitor

REQUIRED MEDICAL INFORMATION

1. Diagnosis of cryopyrin-associated periodic syndrome 2. Diagnosis of recurrent pericarditis 3. Diagnosis of deficiency of Interleukin-1 Receptor Antagonist (DIRA)

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

BvsD Determination

PART B PREREQUISITE

N/A

ATOPIC DERM

MEDICATION(S)

ADBRY, CIBINQO

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

Used in combination with another biologic (e.g., Dupixent, Xolair, Humira, Entyvio, and Otezla, etc.)

REQUIRED MEDICAL INFORMATION

1. Diagnosis of moderate to severe atopic dermatitis (AD)
 - A. Greater than or equal to 10 percent body surface area coverage
 - B. Failure of two of the following:
 - i. Topical corticosteroid (e.g., fluocinonide, clobetasol, etc.)
 - ii. Topical calcineurin inhibitor (eg. tacrolimus ointment 0.1%)
 - iii. Phototherapy
 - iv. Oral immunomodulator (azathioprine, cyclosporine, or mycophenolate)
 - v. Topical PDE-4 (Eucrisa)
2. For reauthorization of Adbry
 - A. a) For patients less than 100kg who achieved clear or almost clear skin, every 4 week dosing has been tried

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

AUGTYRO

MEDICATION(S)

AUGTYRO

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of locally advanced or metastatic ROS1-positive non-small cell lung cancer (NSCLC) or NTRK gene fusion-positive solid tumors
2. Contraindication, intolerance, or failure of Rozlytrek

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

AUSTEDO

MEDICATION(S)

AUSTEDO, AUSTEDO XR, AUSTEDO XR PATIENT TITRATION

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of chorea associated with Huntington's disease 2. Dx of tardive dyskinesia

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

AYVAKIT

MEDICATION(S)

AYVAKIT

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1)Diagnosis of:

- a)Unresectable or metastatic GIST that is platelet-derived growth factor receptor alpha (PDGFRA) exon 18 mutation positive including PDGFRA D842V mutations.
- b)Advanced Systemic Mastocytosis: AdvSM includes patients with aggressive systemic mastocytosis (ASM), systemic mastocytosis with an associated hematological neoplasm (SM-AHN), and mast cell leukemia (MCL).
- c)Indolent Systemic Mastocytosis

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

BALVERSA

MEDICATION(S)

BALVERSA

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Diagnosis of locally advanced or metastatic urothelial carcinoma
2. Confirmed fibroblast growth factor receptor (FGFR3) genetic alteration
3. Progression on or after at least one line of prior systemic therapy

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

BANZEL

MEDICATION(S)

RUFINAMIDE

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Pt has inadequate seizure control despite treatment with at least ONE anti-epileptic drug

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

BENLYSTA

MEDICATION(S)

BENLYSTA 200 MG/ML SOLN A-INJ, BENLYSTA 200 MG/ML SOLN PRSYR

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Diagnosis of lupus nephritis or systemic lupus erythematosus
2. Pt is currently autoantibody positive,
3. Pt has current active disease,
4. Pt has previous treatment with at least TWO of the following:
 - a. Corticosteroids, b. Antimalarials, c. Immunosuppressives,
5. Pt will continue to receive concomitant standard treatment with at least ONE of the following:
 - a. Corticosteroids, b. Antimalarials, c. Immunosuppressives

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

BvsD Determination

PART B PREREQUISITE

N/A

BESREMI

MEDICATION(S)

BESREMI

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of Polycythemia Vera 2. Inadequate response or intolerance to hydroxyurea

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Initial: 6 months Reauth: Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

BOSULIF

MEDICATION(S)

BOSULIF

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Pt is diagnosed with Philadelphia chromosome positive (Ph+) CML a. Pt's CML is newly diagnosed in chronic phase b. Pt's CML is in chronic phase, accelerated phase, or blast phase, 2. For CML in chronic phase, accelerated phase, or blast phase: a. Pt has previous failure or intolerance to imatinib

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

BRAFTOVI

MEDICATION(S)

BRAFTOVI 75 MG CAP, MEKTOVI

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of BRAF V600E or V600K mutation-positive unresectable or metastatic melanoma a. Will be used as combination therapy with encorafenib [Braftovi] and binimetinib [Mektovi] 2. For encorafenib [Braftovi], dx of BRAF V600E mutation-positive metastatic colorectal cancer a. Will be used as combination therapy with cetuximab [Erbix] 3. Dx of BRAF V600E mutation-positive metastatic non-small cell lung cancer (NSCLC) a. Will be used as combination therapy with encorafenib [Braftovi] and binimetinib [Mektovi]

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

BRONCHITOL

MEDICATION(S)

BRONCHITOL, BRONCHITOL TOLERANCE TEST

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of cystic fibrosis 2. Trial of both Hypersal and sodium chloride (3% or 7%)

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

BRUKINSA

MEDICATION(S)

BRUKINSA

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

- 1)Diagnosis of
 - a)Mantle cell lymphoma (MCL)
 - b)Marginal zone lymphoma (MZL)
 - c)Waldenstrom macroglobulinemia
 - d)Chronic lymphocytic leukemia (CLL)/ Small lymphocytic lymphoma (SLL)
 - e)Relapsed or refractory follicular lymphoma (FL)
- 2)Patient has received at least one prior therapy for MCL
- 3)Patient has been previously treated with anti-CD20-based regimen for MZL
- 4)Patient has received two or more lines of therapy for FL and Brukinsa will be used in combination with obinutuzumab

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

CABLIVI

MEDICATION(S)

CABLIVI

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

A. Initial authorization 1. Diagnosis of acquired thrombotic thrombocytopenic purpura (aTTP) 2. Started inpatient in combination with plasma exchange B. Reauthorization 1. Signs of persistent underlying disease (e.g., suppressed ADAMTS13 concentrations) 2. Demonstrated a positive response to therapy by a clinically significant increase in platelet count, reduction in neurological symptoms, or improvement in organ-damage markers

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

3 months

OTHER CRITERIA

Max Duration of therapy 58 days

PART B PREREQUISITE

N/A

CABOMETYX

MEDICATION(S)

CABOMETYX

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of hepatocellular carcinoma a. Previous use of sorafenib 2. Dx of advanced renal cell carcinoma 3. Dx of locally advanced or metastatic differentiated thyroid cancer (DTC) a. progression following prior VEGFR-targeted therapy b. patient is radioactive iodine-refractor or ineligible

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

CALQUENCE

MEDICATION(S)

CALQUENCE

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of mantle cell lymphoma a. Pt has been treated with at least one prior therapy 2. Dx of chronic lymphocytic leukemia 3. Dx of small lymphocytic lymphoma

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

CAPRELSA

MEDICATION(S)

CAPRELSA

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx symptomatic or progressive medullary thyroid cancer with unresectable locally advanced or metastatic disease

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

CARBAGLU

MEDICATION(S)

CARGLUMIC ACID, SAPROPTERIN DIHYDROCHLORIDE

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. For carglumic acid: hyperammonemia due to N-acetylglutamate synthase deficiency 2. For sapropterin: hyperphenylalaninemia due to tetra hydrobiopterin- (BH4-) responsive Phenylketonuria

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

CHOLBAM

MEDICATION(S)

CHOLBAM

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Initial: a. Pt has abnormal results from a urinary bile acids analysis by FAB-MS and neurologic exam,
2. Reauth: a. Patient has experienced improvement in ALT/AST values, bilirubin values, and/or weight

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Initial: Plan year Reauth: Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

COMETRIQ

MEDICATION(S)

COMETRIQ (100 MG DAILY DOSE), COMETRIQ (140 MG DAILY DOSE), COMETRIQ (60 MG DAILY DOSE)

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of a. Progressive, metastatic medullary thyroid cancer (MMTC) 2. Max daily dose of 140 mg/day

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

COPIKTRA

MEDICATION(S)

COPIKTRA

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

Previous use of 2 prior therapies for indication

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

CRINONE

MEDICATION(S)

CRINONE 4 % GEL

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

A. Diagnosis of secondary amenorrhea

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

CYSTAGON

MEDICATION(S)

CYSTAGON

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Diagnosis of nephropathic cystinosis 2. Elevated baseline WBC cysteine levels greater than 2 nmol per 1/2 cystine/mg protein 3. CTNS gene mutation 4. Clinical symptoms of an electrolyte imbalance and polyuria

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

CYSTARAN

MEDICATION(S)

CYSTADROPS, CYSTARAN

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Initial: a. Pt has corneal cysteine accumulation that has been confirmed by slit-lamp photography,

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

CYSTIC FIBROSIS

MEDICATION(S)

CAYSTON, TOBRAMYCIN 300 MG/5ML NEBU SOLN

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Pt has had at least ONE positive culture for Pseudomonas aeruginosa 2. If request not for generic tobramycin: Previous trial on generic tobramycin

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

BvsD determination for tobramycin only, no BvD for Cayston

PART B PREREQUISITE

N/A

DANZITEN

MEDICATION(S)

DANZITEN

PENDING CMS APPROVAL

DAURISMO

MEDICATION(S)

DAURISMO

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Equal to or greater than 75 years or has comorbidity preventing use of intensive induction chemotherapy. 2. Be given in combination with low-dose cytarabine

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

DIACOMIT

MEDICATION(S)

DIACOMIT

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of Dravet syndrome 2. Previous use of clobazam and valproic acid 3. To be used in combination with clobazam

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

DOPTELET

MEDICATION(S)

DOPTELET

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Platelet count less than 50,000 2. Pt is scheduled for a procedure where there is a bleeding risk 3. Doptelet will be used for 5 days starting 10 to 13 days prior to the procedure and discontinued 5 to 8 days prior to the procedure 4. For chronic immune thrombocytopenia, has the patient had an insufficient response to a previous treatment (e.g. corticosteroid, immune globulin)

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

DRONABINOL

MEDICATION(S)

DRONABINOL

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. For HIV-associated wasting syndrome OR cancer-associated anorexia dx: a. Pt has previous trial on megestrol, 2. For CINV dx: a. Pt has previous trial on olanzapine

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

DUPIXENT

MEDICATION(S)

DUPIXENT 300 MG/2ML SOLN A-INJ, DUPIXENT 300 MG/2ML SOLN PRSYR

PA INDICATION INDICATOR

2 - Some FDA-Approved Indications Only

OFF LABEL USES

N/A

EXCLUSION CRITERIA

1. Use in combination with any other biologic (e.g., Dupixent, Xolair, Humira, Entyvio, and Otezla, etc.)
2. Use in combination with Ohtuvayre

REQUIRED MEDICAL INFORMATION

1. Diagnosis of corticosteroid-dependent asthma (Please note, Dupixent is only covered for corticosteroid-dependent asthma, chronic obstructive pulmonary disease, and prurigo nodularis)
 - A. Diagnosis of corticosteroid-dependent asthma evidenced by members requirement of at least 3 months of chronic oral corticosteroids
 - B. 3 month failure of at least TWO of the following:
 - i. ICS + SABA
 - ii. ICS/LABA
 - iii. Spiriva or Trelegy
 - iv. LTRA
 - C. Two provider/ER/Hospital admissions for asthma exacerbations within the last 12 months
2. Diagnosis of prurigo nodularis (PN)
 - A. Pt has a PP-NRS score of 7 or more
 - B. Pt has a history of failing a 2-week course of topical corticosteroids
3. Diagnosis of Chronic Obstructive Pulmonary Disease (COPD)
 - A. Diagnosis confirmed by post-bronchodilator FEV1/FVC less than 0.7 on spirometry
 - B. Blood eosinophil count (BEC) greater than or equal to 300 cells/mcL within last 3 months
 - C. Chronic bronchitis, defined as chronic productive cough for greater than or equal to 3 months in the past year, in absence of other known causes of chronic cough
 - D. 2 ER/hospital admissions for COPD exacerbations within the last year despite at least 3 months treatment with inhaled LABA + LAMA + ICS triple therapy (or LABA + LAMA if ICS is contraindicated)

AGE RESTRICTION

For PN: 18 years of age or older

PRESCRIBER RESTRICTION

For corticosteroid dependent asthma: pulmonologist or allergist

For COPD: pulmonologist

For PN: dermatologist, allergist, or immunologist

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

EMGALITY

MEDICATION(S)

EMGALITY, EMGALITY (300 MG DOSE)

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Diagnosis of episodic cluster headache 2. Diagnosis of Episodic Migraine (4-14 migraine days per month) or Chronic Migraine (greater than 14 migraine days per month) A. Member has tried Ajovy
Reauthorization Criteria: 1. Patient has had a reduction in the number of migraine days per month

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

ENDARI

MEDICATION(S)

L-GLUTAMINE 5 GM PACKET

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

1. Renal insufficiency, 2. Uncontrolled liver disease

REQUIRED MEDICAL INFORMATION

1. Dx of Sickle Cell Disease (SCD), 2. Tx to prevent acute complications of sickle cell disease, 3. Previous use, concurrent use, or inability to use generic hydroxyurea, 4. Reauthorization: reduction in the number of acute complications (i.e blood transfusions, sickle cell crisis, hospitalizations) of sickle cell disease since initiating therapy

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

ENDOTHELIN ANTAGONISTS

MEDICATION(S)

ADEMPAS, AMBRISENTAN, BOSENTAN, OPSUMIT

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. PAH:

- a) Exclusion of all secondary causes of pulmonary hypertension
- b) Must be dx with PAH with WHO class II, III, or IV
- c) For Opsumit or Adempas: Previous trial of ambrisentan or bosentan

2. CTPH (for Adempas):

- a) Pt has failed endarterectomy OR
- b) Pt considered inoperable for pulmonary endarterectomy

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

ENSPRYNG

MEDICATION(S)

ENSPRYNG

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

1. Use of another biologic tx for NMOSD

REQUIRED MEDICAL INFORMATION

1. Dx of neuromyelitis optica spectrum disorder (NMOSD) with one of the following:
 - A. Idiopathic single or recurrent events of longitudinally extensive myelitis (3 or more vertebral segment spinal cord MRI lesion)
 - B. Optic neuritis, single, recurrent or simultaneous bilateral
2. Positive for anti-aquaporin-4 (AQP4) antibody,
3. Reauthorization:
 - A. Patient is continuing to receive benefit from treatment
4. Chart notes required

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Initial: Plan year Reauth: Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

EPCLUSA

MEDICATION(S)

SOFOSBUVIR-VELPATASVIR

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Chart notes documenting genotype,

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

12 Weeks

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

EPIDIOLEX

MEDICATION(S)

EPIDIOLEX

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

Previous use of two alternative antiepileptic medications and used in combination with another antiepileptic

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

ERIVEDGE

MEDICATION(S)

ERIVEDGE, ODOMZO

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Pt has recurring lesions after radiation therapy OR radiation therapy is contraindicated or inappropriate, 2. Pt has recurring lesions after surgical excision OR surgery is contraindicated or inappropriate

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

ERLEADA

MEDICATION(S)

ERLEADA

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

- 1)Diagnosis of metastatic castration sensitive prostate cancer (mCSPC)
 - a)Patient has tried and failed abiraterone 250mg tablets AND Xtandi
- 2)Diagnosis of non-metastatic castration-resistant prostate cancer (nmCRPC)
 - a)Patient has PSA doubling time of less than 10 months
 - b)Patient has failed Xtandi

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

ESBRIET

MEDICATION(S)

PIRFENIDONE

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx: a. Idiopathic pulmonary fibrosis (IPF) 2. Dx has been confirmed via high-resolution computed tomography scan and/or lung biopsy, 3. Forced Vital Capacity (FVC) greater than 50 percent predicted value 4. Carbon monoxide diffusing capacity (DLCO) greater than 30 percent predicted value 5. Liver function tests (ALT, AST, and bilirubin) completed prior to initiating therapy 6. Reauth: a. A repeat liver function test has been performed after 3 months of therapy has been completed

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Initial: 3 mo, Reauth: Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

ETHACRYNIC ACID

MEDICATION(S)

ETHACRYNIC ACID

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Pt needs diuresis for any of the following: a. Edema associated with congestive heart failure, b. Edema associated with cirrhosis of the liver, c. Edema associates with renal disease. d. Short-term management of ascites due to malignancy, idiopathic edema, or lymphedema. 2. Pt has previous trial and failure on a loop diuretic or thiazide diuretic

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

FARESTON

MEDICATION(S)

TOREMIFENE CITRATE

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

1. Pt has congenital or acquired QT prolongation, 2. Pt has uncorrected hypokalemia, 3. Pt has uncorrected hypomagnesemia

REQUIRED MEDICAL INFORMATION

1. Pt has previous trial and failure or contraindication to tamoxifen therapy, 2. Pt has previous trial and failure or contraindication to aromatase inhibitor therapy

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

FASENRA

MEDICATION(S)

FASENRA, FASENRA PEN

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

1. Concurrent therapy with another biologic medication

REQUIRED MEDICAL INFORMATION

1. Dx of Severe Asthma

a. Patient has tried at least 3 months on any of the following combinations:

- i. ICS/LABA
- ii. ICS/LTRA
- iii. ICS/LAMA
- iv. ICS/LABA/LAMA (Trelegy)

b. Two exacerbations requiring the use of oral corticosteroids in the previous 12 months OR one exacerbation that led to a hospitalization in the previous 12 months

c. Peripheral blood eosinophil level greater than 150 cells/mcL

2. Dx of eosinophilic granulomatosis with polyangiitis (EGPA)

a. Patient must have TWO of the following disease characteristics

i. Biopsy showing histopathological evidence of eosinophilic vasculitis, perivascular eosinophilic infiltration, or eosinophil-rich granulomatous inflammation

- ii. Neuropathy
- iii. Pulmonary infiltrates, non-fixed
- iv. Sino-nasal abnormality
- v. Cardiomyopathy (established by echocardiography or MRI)
- vi. Glomerulonephritis (hematuria, red cell casts, proteinuria)
- vii. Alveolar hemorrhage
- viii. Palpable purpura
- ix. Anti-neutrophil cytoplasmic anti-body (ANCA) positive

c. Patient has a history of eosinophil level greater than 100 cells/mcL or blood eosinophil level

greater than 10%

d. Patient has a history of relapse or refractory disease despite current use of oral glucocorticoids unless contraindicated or not tolerated

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

For EGPA: allergist, immunologist, rheumatologist, or pulmonologist

COVERAGE DURATION

Plan Year

OTHER CRITERIA

BvsD Determination

PART B PREREQUISITE

N/A

FENTANYL

MEDICATION(S)

FENTANYL CITRATE 1200 MCG LOZ HANDLE, FENTANYL CITRATE 1600 MCG LOZ HANDLE, FENTANYL CITRATE 200 MCG LOZ HANDLE, FENTANYL CITRATE 400 MCG LOZ HANDLE, FENTANYL CITRATE 600 MCG LOZ HANDLE, FENTANYL CITRATE 800 MCG LOZ HANDLE

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

Non-cancer pain use

REQUIRED MEDICAL INFORMATION

1. Must have documented maintenance therapy with a long-acting opioid, 2. Documented failure with a short-acting opioid for breakthrough pain

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

6 months

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

FENTANYL PATCH

MEDICATION(S)

FENTANYL 100 MCG/HR PATCH 72HR, FENTANYL 12 MCG/HR PATCH 72HR, FENTANYL 25 MCG/HR PATCH 72HR, FENTANYL 50 MCG/HR PATCH 72HR, FENTANYL 75 MCG/HR PATCH 72HR

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

- 1.Fentanyl used for acute intermittent pain, post-operative pain, or mild pain
- 2.Fentanyl used in combination with a benzodiazepine or other central nervous system (CNS) depressant

REQUIRED MEDICAL INFORMATION

- 1.Patient has severe pain requiring long-term, around the clock pain management for which alternative treatment options have been inadequate
- 2.Provider is knowledgeable in the use of potent opioids for chronic pain management
- 3.Patient is opioid tolerant, having been established on the equivalent of 60 milligrams of morphine or more per day for at least a week prior to being prescribed fentanyl patches
- 4.Patient will be monitored for sedation and respiratory depression
- 5.Patient will be monitored for opioid addiction, abuse, and misuse
- 6.Appropriate use and availability of naloxone has been discussed with the patient
- 7.If requesting every 48-hour dosing, every 72-hour dosing has been tried
8. Patient has tried a dose reduction within the last 6 months

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

6 months

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

FILSUVEZ

MEDICATION(S)

FILSUVEZ

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

Combination use with Vyjuvek

REQUIRED MEDICAL INFORMATION

1. Dx of dystrophic or junctional epidermolysis bullosa
2. Prescribing physician is a dermatologist with experience in treating epidermolysis bullosa and collaborated with a wound healing specialist
3. Reauthorization:
 - a. Documentation of a response as evidenced by wound healing

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

Dermatologist

COVERAGE DURATION

Initial: 6 months

Reauth: Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

FINTEPLA

MEDICATION(S)

FINTEPLA

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of Dravet Syndrome or Lennox Gaustaut Syndrome
2. Previous use of two of topiramate, valproic acid, or clobazam

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

FIRAZYR

MEDICATION(S)

ICATIBANT ACETATE

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. For HAE type I and II and acquired angioedema: a. Dx has been verified by low C1-INH and/or low C1-INH function levels
2. For HAE with normal C1-INH:
 - a. Pt has failed a trial with high-dose non-sedating antihistamines such as cetirizine, desloratadine, or levocetirizine for at least 30 days to rule-out idiopathic angioedema

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

1. Initial: 6 months, 2. Reauth: Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

FIRDAPSE

MEDICATION(S)

FIRDAPSE

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

I. Initial authorization

A. P/Q-type voltage-gated calcium channel antibodies OR Repetitive nerve stimulation consistent with LEMS

B. Screening for cancer related to LEMS

C. Experiences moderate to severe weakness interfering with function

D. Documentation of quantitative myasthenia gravis core and subjective global impression score

E. Patient has tried and failed pyridostigmine

II. Reauthorization

A. Improvements in myasthenia gravis core and subjective global impression score

B. Screened 3-6 months after initial screening for malignancies

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Initial: 3 months Reauthorization: 6 months

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

FOTIVDA

MEDICATION(S)

FOTIVDA

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Diagnosis of relapsed or refractory advanced renal cell carcinoma (RCC) 2. Previous failure of two systemic therapies

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

FRUZAQLA

MEDICATION(S)

FRUZAQLA

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of metastatic colorectal cancer (mCRC) 2. Member has previously been treated with fluoropyrimidine-, oxaliplatin-, and irinotecan-based chemotherapy, an anti-VEGF therapy, and, if RAS wild-type and medically appropriate, an anti-EGFR therapy 3. Member has failed, contraindication, or intolerance to Lonsurf with or without bevacizumab

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

GAVRETO

MEDICATION(S)

GAVRETO

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of non-small cell lung cancer (NSCLC) A. Metastatic NSCLC identified as rearranged during transfection (RET) fusion-positive 2. Dx of thyroid cancer A. Advanced or metastatic RET fusion-positive thyroid cancer refractory radioactive iodine (if appropriate) requiring systemic therapy

AGE RESTRICTION

12 and older

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

GILOTRIF

MEDICATION(S)

GILOTRIF

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of metastatic non-small cell lung cancer (NSCLC) a. The tumor has an epidermal growth factor receptor (EGFR) exon 19 deletion or exon 21 (L858R) substitution mutation 2. Dx of metastatic squamous non-small cell lung cancer (NSCLC) a. Progression after platinum-based chemotherapy

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

GLEOSTINE

MEDICATION(S)

GLEOSTINE

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of primary and metastatic brain tumors following appropriate surgical and/or radiotherapeutic procedures
2. Dx of Hodgkin's lymphoma A. Disease has progressed following initial chemotherapy

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

GLP 1

MEDICATION(S)

MOUNJARO, TRULICITY

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

Type 1 diabetes mellitus

REQUIRED MEDICAL INFORMATION

- 1) Diagnosis of Type 2 diabetes mellitus (T2DM) AND
- 2) Patient has tried and failed a 90-day trial of a sodium-glucose cotransporter-2 (SGLT2) inhibitor (e.g., Farxiga, Glyxambi, Invokamet, Segluromet, Synjardy, Xigduo, etc.) or a dipeptidyl peptidase IV (DPP4) inhibitor (e.g., alogliptin, Januvia, Janumet, Onglyza, Tradjenta, etc.) as evidenced by an A1c greater than or equal to 7% in last 3 months
 - a) OR has ONE of the following
 - i) Reduced renal function (eGFR 45mL/min/m² or less)
 - ii) Urinary frequency due to BPH, LUTS, bladder spasm, etc.
 - iii) Recurrent genital fungal infection
 - iv) Recurrent urinary tract infection
 - v) Intolerance to an SGLT2/DDP4 inhibitor
 - b) AND the patient has an A1c greater than or equal to 7% in the last 3 months

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

GROWTH HORMONE

MEDICATION(S)

GENOTROPIN, GENOTROPIN MINIQUICK, OMNITROPE

PA INDICATION INDICATOR

4 - All FDA-Approved Indications, Some Medically-Accepted Indications

OFF LABEL USES

1. All medically accepted indications not otherwise excluded from Part D

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1)Adult Growth Hormone Deficiency

a)Patient has failed two of the following stimulation tests:

i)Arginine, Clonidine, Glucagon, Insulin, Levodopa, Gonadotropin

ii)OR patient has at least two other pituitary hormone deficiencies in addition to low insulin-like growth factor 1 measurement below the age appropriate level (e.g. below 2.5th percentile or Z-score less than -2

iii)OR the patient has already been established on growth hormone at a pediatric age (under 18)

2)Pediatric growth hormone deficiency (Ped GHD) or idiopathic short stature (ISS)

3)Chronic Renal Insufficiency:

a)Meet ALL of the following:

i)Pt dxed with CRI AND has not yet received renal transplant,

ii)Existing metabolic disorders have been corrected,

iii)Ht more than 2 SD below the population mean OR less than 3rd percentile,

iv)Height velocity less than 4cm/yr or less than 10th percentile of normal for age and gender,

4)Turner Syndrome:

a)Meet ALL of the following:

i)Dx of TS confirmed by blood karyotype or fibroblast studies,

ii)Ht of female pt plotted on TS-specific growth curve AND pt is less than 5th percentile of normal growth curve for girls,

5)Prader-Willi Syndrome:

a)Meet ALL of the following:

i)Dx of PWS confirmed by appropriate genetic testing,

- ii)Ht more than 2 SD below the pop mean OR less than 3rd percentile,
- iii)Ht velocity less than 3cm/yr or less than 10th percentile of normal for age and gender,
- 6)Small for Gestational Age:
 - a)Meet ALL of the following:
 - i)Dx of SGA as defined as one of the following:
 - (1)Birth weight of less than 2,500g at gestational age of greater than 37 weeks,
 - (2)OR birth weight or length less than 3rd percentile for gestational age,
 - ii)Pt has failed to catch up in ht by 2 yo,

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

- 1)HIV wasting: 48 wks LIFETIME,
- 1)All other indications: Plan Year

OTHER CRITERIA

- 7)AIDS-Related Wasting:
 - a)Meet ALL of the following:
 - i)Involuntary weight loss of more than 10% pre-illness body weight or a BMI less than 20,
 - ii)Failure to respond to dronabinol (Marinol) OR megestrol acetate (Megace),
 - iii)Chronic diarrhea (defined as more than 3 loose stools/day for more than 30 days) OR Chronic weakness and documented fever (30 days, intermittent or constant) in the absence of concurrent illness or condition other than HIV infection that would otherwise explain the symptoms.

PART B PREREQUISITE

N/A

HAEGARDA

MEDICATION(S)

HAEGARDA, ORLADEYO

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Initial:

- a. Dx of HAE type I or II has been verified by low C1-INH and/or low C1-INH functional levels
 - i. Pt has a history of facial, laryngeal, and/or gastrointestinal HAE attacks
- b. Dx of HAE with normal C1-INH:
 - i. Pt has failed a trial with high-dose non-sedating antihistamines such as cetirizine, desloratadine, or levocetirizine for at least 30 days to rule-out idiopathic angioedema

2. Reauthorization:

- a. Pt had a significant decrease in the frequency of attacks per month or had a significant decrease in the severity or duration of attacks

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

HARVONI

MEDICATION(S)

LEDIPASVIR-SOFOSBUVIR

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Chart notes showing genotype

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Maximum 24 weeks

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

HEMADY

MEDICATION(S)

HEMADY

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Diagnosis of Multiple Myeloma (MM) 2. Previous use of generic dexamethasone tablets

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

HEPATITIS C

MEDICATION(S)

PEGASYS

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. For Hep B dx: a. Pre-treatment HBV DNA levels are greater than 20,000 IU/ml, b. Must be used as monotherapy 2. Hep C dx

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

48 weeks

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

HETLIOZ

MEDICATION(S)

HETLIOZ LQ, TASIMELTEON

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. For Non-24 Hour Sleep-Wake Disorder: a. Pt is totally blind without light perception 2. For Nighttime Sleep Disturbances in Smith-Magenis Syndrome (SMS) a. Diagnosis been confirmed with genetic testing

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

Sleep specialist or Neurologist

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

HUMIRA

MEDICATION(S)

AMJEVITA 20 MG/0.2ML SOLN PRSYR, AMJEVITA 40 MG/0.4ML SOLN A-INJ, AMJEVITA 40 MG/0.4ML SOLN PRSYR, AMJEVITA 80 MG/0.8ML SOLN A-INJ, AMJEVITA-PED 15KG TO <30KG, HADLIMA, HADLIMA PUSHTOUCH

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

1. Combination therapy with another biologic medication, JAK Inhibitor, or Otezla

REQUIRED MEDICAL INFORMATION

1. Dx of AS, UC, or CD 2. Dx of RA, JIA, PsA dx: Pt has failed at least three months therapy on at least ONE of the following: a. methotrexate, b. leflunomide, c. hydroxychloroquine, d. sulfasalazine, OR contraindication to use (Clinical diagnosis of alcohol use disorder, alcoholic liver disease or other chronic liver disease, Pregnancy, breastfeeding, or currently planning pregnancy, Blood dyscrasias (e.g., thrombocytopenia, leukopenia, significant anemia), Elevated liver transaminases, Hypersensitivity, or history of intolerance or adverse event, Interstitial pneumonitis or clinically significant pulmonary fibrosis, Myelodysplasia, Renal impairment 3. Dx of hidradenitis suppurativa: a. Pt has failed therapy or had an inadequate response to a treatment of oral antibiotics b. Pt has lesions present in at least TWO distinct anatomical areas, one of which is Hurley Stage II or III 4. Dx of noninfectious uveitis: a. Pt has previous failure on corticosteroids 5. Dx of plaque psoriasis: Pt has failed therapy with at least ONE of the following: a. methotrexate, b. cyclosporine, c. acitretin

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

HYFTOR

MEDICATION(S)

HYFTOR

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of facial angiofibroma associated with tuberous sclerosis 2. Member's facial angiofibroma cause functional impairment or symptoms such as bleeding or pain

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

12 weeks No reauthorization

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

IBRANCE

MEDICATION(S)

IBRANCE

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1)Dx of hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative advanced or metastatic breast cancer

a)Ibrance used as initial therapy:

i)Used in combination with an aromatase inhibitor,

b)Ibrance used after endocrine-based therapy:

i)Used in combination with fulvestrant

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

ICLUSIG

MEDICATION(S)

ICLUSIG

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Chronic Phase CML dx
 - a. Resistance or intolerance to at least two prior kinase inhibitors.
2. Accelerated or Blast phase CML OR Ph+ ALL
 - a. No other TKI is indicated
3. T315I-positive CML or T315I-positive ALL
4. Newly diagnosed Ph+ ALL

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

IDHIFA

MEDICATION(S)

IDHIFA

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Cancer has an isocitrate dehydrogenase-2 (IDH2) mutation as detected by a FDA-approved test,
2. Pt has relapsed or is refractory to one or more prior anticancer regimens

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

IMBRUVICA

MEDICATION(S)

IMBRUVICA 140 MG CAP, IMBRUVICA 140 MG TAB, IMBRUVICA 280 MG TAB, IMBRUVICA 420 MG TAB, IMBRUVICA 70 MG CAP, IMBRUVICA 70 MG/ML SUSPENSION

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of CLL/SLL 2. Dx of CLL/SLL w/ 17p deletion 3. Dx of Waldenstrom's Macroglobinemia 4. Dx of cGVHD: a. Member has failed at least one prior systemic therapy b. Prescribing physician is a specialist in transplant

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

IMKELDI

MEDICATION(S)

IMKELDI

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Pt is unable to swallow tablets

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

IMPAVIDO

MEDICATION(S)

IMPAVIDO

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Diagnosis of leishmaniasis (visceral, cutaneous, or mucosal) confirmed by parasite in a clinical specimen 2. Previous use of ketoconazole, fluconazole, paromomycin, or amphotericin 3. Weight 30 kg or greater

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

28 days

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

INCRELEX

MEDICATION(S)

INCRELEX

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Height standard deviation score of less than -3 based on age and gender, 2. Basal IGF-1 standard deviation score of less than -3 based on age and gender, 3. Normal or elevated growth hormone levels, 4. Pt must have open epiphyses, 5. Gh stimulation test of greater than 10 mcg/L.

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

INLYTA

MEDICATION(S)

INLYTA

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of renal cell carcinoma 2. For first line treatment in combination with pembrolizumab (Keytruda) or avelumab (Bavencio) 3. Monotherapy as a second line treatment after previous use of one prior systemic therapy

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

INQOVI

MEDICATION(S)

INQOVI

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of myelodysplastic syndrome

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

INREBIC

MEDICATION(S)

INREBIC

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

Intolerance, failure, or previous use of ruxolitinib

REQUIRED MEDICAL INFORMATION

A. Diagnosis of intermediate or high-risk primary or secondary myelofibrosis B. Platelet count greater than or equal to $50 \times 10^9/L$

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

IRESSA

MEDICATION(S)

GEFITINIB

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Pt has ONE of the following: a. EGFR exon 19 deletion, b. EGFR exon 21 deletion, 2. Gefitinib is used as first-line therapy

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

ISTURISA

MEDICATION(S)

ISTURISA 1 MG TAB, ISTURISA 5 MG TAB

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of Cushings Disease a. Baseline mean urinary free cortisol (UFC) level at least 1.5x the upper limit of normal measured over three 24 hour measurements (ULN = 50 micrograms/24 hours or 145 nmol/24 hours) b. Symptoms of Cushings Disease (e.g diabetes, central obesity, moon face, buffalo hump, osteoporosis, muscle wasting, hypertension, depression, anxiety) c. Failure of pituitary surgery or contraindication to pituitary surgery d. Intolerance to pasireotide (Signifor) g. Exclusion of other causes of Cushings Syndrome (aside from Cushings Disease which is specifically caused by a pituitary adenoma) 2. For reauthorization: a. Recent UFC level showing improvement (less than 48 weeks of treatment) or is within normal limits (after 48 weeks of treatment) b. Symptom improvement of Cushings Disease

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

ITOVEBI

MEDICATION(S)

ITOVEBI

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

1. Disease progression on or following other PI3k inhibitors

REQUIRED MEDICAL INFORMATION

1. Pt must have advanced or metastatic HR-positive, HER2-negative breast cancer with documented PIK3CA mutation via specified testing
2. Pt must have disease progression following at least one line of endocrine therapy in the metastatic setting or recurrence on or within 12 months of completing adjuvant therapy
3. Itovebi will be given in combination with Ibrance (palbociclib) and fulvestrant as first-line therapy

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

IVIG

MEDICATION(S)

BIVIGAM, GAMMAGARD, GAMMAGARD S/D LESS IGA, GAMMAKED 1 GM/10ML SOLUTION, GAMMAPLEX, GAMUNEX-C 1 GM/10ML SOLUTION, OCTAGAM 1 GM/20ML SOLUTION, OCTAGAM 10 GM/100ML SOLUTION, OCTAGAM 10 GM/200ML SOLUTION, OCTAGAM 2 GM/20ML SOLUTION, OCTAGAM 2.5 GM/50ML SOLUTION, OCTAGAM 25 GM/500ML SOLUTION, OCTAGAM 30 GM/300ML SOLUTION, OCTAGAM 5 GM/100ML SOLUTION, PRIVIGEN

PA INDICATION INDICATOR

3 - All Medically-Accepted Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

- 1) For all non-preferred IVIG products (ie Bivigam, Gammaked, Octagam, etc)
- a) Must try TWO of the following: Gammagard, Gamunex, Hizentra, and Privigen OR
 - b) Documented medical reason why preferred agents cannot be used

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

BvD Determination

PART B PREREQUISITE

YES

IWILFIN

MEDICATION(S)

IWILFIN

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of high risk neuroblastoma (HRNB) in patients who have demonstrated at least a partial response to prior multiagent, multimodality therapy including anti-GD2 immunotherapy 2. Maximum duration of 2 years of therapy

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

IXCHIQ

MEDICATION(S)

IXCHIQ

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

Confirmation that the patient is at increased risk of exposure to chikungunya virus

AGE RESTRICTION

18 years of age or older

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

1 month

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

JADENU

MEDICATION(S)

DEFERASIROX 180 MG PACKET, DEFERASIROX 250 MG TAB SOL, DEFERASIROX 360 MG PACKET, DEFERASIROX 500 MG TAB SOL, DEFERASIROX 90 MG PACKET, DEFERASIROX GRANULES

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. For blood transfusion dx: a. Pt's serum ferritin is greater than 1000 mcg/L, b. Pt has failed subcutaneous deferoxamine through an inability to achieve desired goals of therapy or been intolerant to therapy, 2. For non-transfusion-dependent thalassemia dx: a. Pt's liver iron concentration is at least 5 mg Fe per gram of dry weight, b. Pt's serum ferritin is greater than 300 mcg/L, c. Pt has failed subcutaneous deferoxamine through an inability to achieve desired goals of therapy or been intolerant to therapy

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

Part B before Part D step therapy

PART B PREREQUISITE

YES

JAKAFI

MEDICATION(S)

JAKAFI

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Diagnosis of: a. Intermediate or high-risk myelofibrosis, including primary myelofibrosis, post-polycythemia vera myelofibrosis, or post-essential thrombocythemia myelofibrosis b. Polycythemia vera c. Acute Graft-versus-host disease d. Chronic Graft-versus-host disease 2. For myelofibrosis or polycythemia vera, must have at least ONE of the following: a. Pt has enlarged spleen shown by MRI or CT, b. Pt has palpable splenomegaly 3. For myelofibrosis or polycythemia vera, platelet count greater than or equal to $50 \times 10^9/L$ 4. For the treatment of acute graft-versus-host disease patient has previously failed trial of corticosteroids 5. For the treatment of chronic graft-versus-host disease patient has previously failed one or two lines of systemic therapy

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

JAYPIRCA

MEDICATION(S)

JAYPIRCA

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Diagnosis of relapsed/refractory MCL

a. Previous treatment with at least two lines of systemic therapy, including a BTK inhibitor

2. Diagnosis of chronic lymphocytic leukemia/small lymphocytic lymphoma (CLL/SLL)

a. Previous treatment with at least two prior lines of therapy including a BTK inhibitor and a BCL-2 inhibitor

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

JYNARQUE

MEDICATION(S)

JYNARQUE

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

eGFR less than 25ml/min

REQUIRED MEDICAL INFORMATION

1. Diagnosis of autosomal dominant polycystic kidney disease
 - a) Diagnosis has been confirmed by radiology

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

Endocrinologist, nephrologist, cardiologist and hepatologist.

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

KALYDECO

MEDICATION(S)

KALYDECO

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Initial: A. Pt genotyped by an FDA-cleared CF mutation test B. Pt have one mutation in the CFTR gene that is responsive to ivacaftor potentiation based on clinical and/or in vitro assay data C. Pt has FEV1 between 40-90% 2. Reauth: A. Pt has been reassessed since starting therapy, B. Pt's FEV1 has increased since starting therapy

AGE RESTRICTION

Tablet: 6 years old or older, Granules: 1 months old to 5 years old

PRESCRIBER RESTRICTION

Specialist in Cystic Fibrosis or Pulmonologist

COVERAGE DURATION

Initial: 3 months Reauth: Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

KERENDIA

MEDICATION(S)

KERENDIA

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Diagnosis of Type 2 Diabetes a. Prior treatment with one SGLT-2 inhibitor (Farxiga or Jardiance) b. Contraindication to SGLT2 inhibitor i. eGFR 45ml/min/m² or less ii. Urinary Frequency due to BPH, LUTS, bladder spasm iii. Recurrent genital fungal infection or recurrent urinary tract infection

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

KEVEYIS

MEDICATION(S)

KEVEYIS

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

1. Hypersensitivity to dichlorphenamide or other sulfonamides 2. Use in combination with high-dose aspirin 3. Severe pulmonary disease 4. Hepatic insufficiency (e.g Child-Pugh B or C)

REQUIRED MEDICAL INFORMATION

1. Dx of primary hyperkalemic periodic paralysis, primary hypokalemic periodic paralysis, or related variants. 2. Dx confirmed by genetic testing, provocative testing, electromyography, or muscle biopsy 3. Previous use or contraindication to acetazolamide 4. Reauthorization: a. improvement in baseline symptoms (e.g. number of attacks per week or month, severity of attacks, duration of attacks, short-form 36 assessment, etc.)

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Initial: 6 months Reauthorization: Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

KISQALI

MEDICATION(S)

KISQALI (200 MG DOSE), KISQALI (400 MG DOSE), KISQALI (600 MG DOSE), KISQALI FEMARA (200 MG DOSE), KISQALI FEMARA (400 MG DOSE), KISQALI FEMARA (600 MG DOSE)

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative advanced or metastatic breast cancer, must meet ONE of the following:
 - a. Pt will receive an aromatase inhibitor in combination with Kisqali as initial endocrine based therapy for advanced or metastatic disease,
 - b. Pt will receive fluevestrant in combination with Kisqali
2. Dx of hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative stage II or stage III early breast cancer at high risk of recurrence
 - a. Pt will receive an aromatase inhibitor in combination with Kisqali

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

KORLYM

MEDICATION(S)

MIFEPRISTONE 300 MG TAB

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

Pt using long-term corticosteroid

REQUIRED MEDICAL INFORMATION

1. Pt has previously failed surgery or chemotherapy to correct Cushing's disease OR is ineligible for surgery,
2. Pt with type II diabetes diagnosis,
3. If pt is female:
 - a. Pt has negative pregnancy test within past 14 days,
 - b. Pt is currently using non-hormonal form of birth control

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

Endocrinologist

COVERAGE DURATION

Initial: 6 months, Reauth: Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

KOSELUGO

MEDICATION(S)

KOSELUGO

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Patient has symptomatic, inoperable plexiform neurofibromas (PN)

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

KYNMOBI

MEDICATION(S)

APOMORPHINE HCL

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of Parkinson's disease A. Experiencing off episodes 2. Currently taking an oral formulation of carbidopa/levodopa or there is documentation of an inability to take an oral formulation of carbidopa/levodopa 3. Previous use of an immediate release or oral disintegrating carbidopa/levodopa as a rescue for off episodes 4. Previous use of at least one of: a. COMT inhibitor (tolcapone, entacapone), b. Dopamine agonist (ropinirole, pramipexole), c. MAO-B inhibitor (selegiline, rasagiline, safinamide) 5. Prescribed in combination with antiemetic therapy (Not a 5HT3)

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

LAMPIT

MEDICATION(S)

LAMPIT

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of Chagas disease (T. cruzi)

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

3 months

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

LAZCLUZE

MEDICATION(S)

LAZCLUZE

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

1. Prior systemic treatment for locally advanced Stage III or metastatic Stage IV disease
2. Symptomatic or previously treated unstable brain metastases

REQUIRED MEDICAL INFORMATION

1. Initial

A. Patient must have confirmed NSCLC that is metastatic or unresectable with EGFR exon 19 deletions or 21 L858R substitution mutations.

B. Patient must receive anticoagulant prophylaxis to prevent VTE events for the first 4 months of treatment.

C. Patients must have an ECOG of 0 or 1

D. Clinical rationale why Tagrisso with /without chemotherapy is not appropriate for use in the patient.

2. Reauthorization

A. Tumor assessment does not show new growth, activity or mutations

AGE RESTRICTION

18 years of age or older

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

6 months

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

LENVIMA

MEDICATION(S)

LENVIMA (10 MG DAILY DOSE), LENVIMA (12 MG DAILY DOSE), LENVIMA (14 MG DAILY DOSE), LENVIMA (18 MG DAILY DOSE), LENVIMA (20 MG DAILY DOSE), LENVIMA (24 MG DAILY DOSE), LENVIMA (4 MG DAILY DOSE), LENVIMA (8 MG DAILY DOSE)

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Renal cell carcinoma dx: a. Will be used in combination with everolimus OR pembrolizumab 2. Thyroid cancer dx: a. Tumor is refractory to treatment with radioactive iodine, b. Used as monotherapy 3. Unresectable hepatocellular carcinoma dx: 4. Endometrial carcinoma dx: a. Will be used in combination with pembrolizumab (Keytruda) b. Does not have microsatellite instability-high or mismatch repair deficiency c. Pt has previously been treated with systemic therapy d. Pt is not a candidate for surgery or radiation

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

LEUKINE

MEDICATION(S)

LEUKINE

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of:

a. To shorten time to neutrophil recovery and to reduce the incidence of severe and life-threatening infections and infections resulting in death following induction chemotherapy in adult patients 55 years and older with acute myeloid leukemia (AML).

i. Pt has trial on BOTH of the following: a. Fulphila, b. Udenyca

b, For the mobilization of hematopoietic progenitor cells into peripheral blood for collection by leukapheresis and autologous transplantation.

c. For the acceleration of myeloid reconstitution following autologous bone marrow or peripheral blood progenitor cell transplantation.

d. For the acceleration of myeloid reconstitution following allogeneic bone marrow transplantation.

e. For treatment of delayed neutrophil recovery or graft failure after autologous or allogeneic bone marrow transplantation.

f. To increase survival in patients acutely exposed to myelosuppressive doses of radiation (Hematopoietic Syndrome of Acute Radiation Syndrome [H-ARS]).

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

BvD Determination

PART B PREREQUISITE

N/A

LIVTENCITY

MEDICATION(S)

LIVTENCITY

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

1. Pt is on any other CMV antivirals

REQUIRED MEDICAL INFORMATION

1. Pt weighs at least 35 kg or more
2. History of HSCT or SOT
3. Diagnosis of post-transplant CMV infection/disease with CMV DNA of more than 2730 IU/mL in whole blood or more than 910 IU/mL in plasma
4. CMV disease refractory to or intolerant of first line antiviral treatment (e.g., ganciclovir, valganciclovir, foscarnet, or cidofovir)

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

8 weeks

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

LONSURF

MEDICATION(S)

LONSURF

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

A. Metastatic colorectal cancer 1. Pt has previous therapy on the following: a. A fluoropyrimidine, b. Oxaliplatin, c. Irinotecan, d. Bevacizumab, 2. If cancer is KRAS wild type, pt has received previous therapy with anti-EGFR therapy B. Metastatic gastric or gastroesophageal junction adenocarcinoma previously treated with at least two prior lines of chemotherapy

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

LORBRENA

MEDICATION(S)

LORBRENA

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

- 1) Dx of metastatic non-small cell lung cancer (NSCLC)
 - a) Tumors are anaplastic lymphoma kinase (ALK)-positive
- 2) Patient has failed/intolerance/contraindication to Alecensa

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

LUCEMYRA

MEDICATION(S)

LOFEXIDINE HCL

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Mitigation of opioid withdrawal symptoms
2. Provider submitted documentation that the patient has been counseled on the risks of taking lofexidine with alcohol, benzodiazepines, and/or barbituates
3. Patient has failed clonidine as part of this opioid discontinuation attempt

AGE RESTRICTION

18 years of age or older

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

14 Days

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

LUMAKRAS

MEDICATION(S)

KRAZATI, LUMAKRAS

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

1. Previous KRAS G12C-targeted therapy (other than current) a. Krazati and Lumakras have similar mechanisms of action and it is not recommended to switch between agents upon progression

REQUIRED MEDICAL INFORMATION

1. Initial Authorization:

A. Lumakras:

- i. Diagnosis of KRAS G12C-mutated locally advanced or metastatic non-small cell lung cancer (NSCLC)
- ii. At least one prior systemic therapy
- iii. Monotherapy

B. Krazati:

- i. Diagnosis of KRAS G12C-mutated locally advanced or metastatic non-small cell lung cancer (NSCLC)
 - a. At least one prior systemic therapy
 - b. Monotherapy
- ii. Diagnosis of KRAS G12C-mutated locally advanced or metastatic colorectal cancer (CRC)
 - a. Received prior treatment with fluoropyrimidine-, oxaliplatin-, and irinotecan-based chemotherapy
 - b. Used in combination with cetuximab

2. Reauthorization:

A. Documentation of stable or improved disease

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Initial: 6 months Reauthorization: Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

LUPKYNIS

MEDICATION(S)

LUPKYNIS

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of lupus nephritis 2. eGFR greater than 45 mL/min/1.73m²

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

LYNPARZA

MEDICATION(S)

LYNPARZA

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

Combination therapy

REQUIRED MEDICAL INFORMATION

1. Ovarian cancer, advanced (BRCA-mutated): a. First-line maintenance therapy for gBRCAm or sBRCAm advanced epithelial ovarian, fallopian tube, or primary peritoneal cancer in adult patients with complete or partial response to first-line platinum-based chemotherapy, OR b. First-line maintenance treatment (in combination with bevacizumab) of advanced epithelial ovarian, fallopian tube, or primary peritoneal cancer in adults who are in complete or partial response to first-line, platinum-based chemotherapy and whose cancer is associated with homologous recombination deficiency (HRD)-positive status, 2. Recurrent ovarian cancer dx: a. Maintenance treatment of recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer in adults who are in complete or partial response to platinum-based chemotherapy. 3. Breast Cancer (BRCA-mutated, HER2-negative) dx: a. For Metastatic Breast Cancer: Pt has previous trial of chemotherapy in neoadjuvant, adjuvant, or metastatic setting i. If HR positive: pt has t/f endocrine therapy or endocrine therapy is inappropriate for pt b. For High Risk Early Breast Cancer: Pt has previous trial of chemotherapy in neoadjuvant, adjuvant, or metastatic setting 4. Pancreatic cancer (BRCA-mutated): a. disease has not progressed on at least 16 weeks of a first-line, platinum-based chemotherapy regimen 5. Prostate cancer (mCRPC): a. Homologous recombination repair (HRR) gene-mutated metastatic castration-resistant prostate cancer in adults who have progressed following prior enzalutamide or abiraterone treatment b. BRCA-mutated (BRCAm) metastatic castration resistant prostate cancer (mCRPC) in combination with abiraterone and prednisone or prednisolone

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

LYTGOBI

MEDICATION(S)

LYTGOBI (12 MG DAILY DOSE), LYTGOBI (16 MG DAILY DOSE), LYTGOBI (20 MG DAILY DOSE)

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Diagnosis of metastatic cholangiocarcinoma 2. Previous treatment for metastatic cholangiocarcinoma with at least 1 line of systemic therapy 3. Fibroblast growth factor receptor 2 (FGFR2) fusion or other rearrangement as detected by an FDA-approved test

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

MAVYRET

MEDICATION(S)

MAVYRET

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

Confirmation of genotype

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

16 Weeks

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

MEKINIST

MEDICATION(S)

MEKINIST

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Pt is BRAF(V600E or V600K) mutation positive, 2. Monotherapy or in combination with Tafinlar 3. For Mekinist oral solution: Member has inability to swallow tablets

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

MIGRANAL

MEDICATION(S)

DIHYDROERGOTAMINE MESYLATE 4 MG/ML SOLUTION

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Pt experiences at least 2 migraines per month,
2. Pt has trial or contraindication to at least TWO of the following:
 - a. Sumatriptan, b. Rizatriptan, c. Zolmitriptan, d. Naratriptan

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

MULPLETA

MEDICATION(S)

MULPLETA

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Platelet count less than 50,000 2. Pt is scheduled for a procedure where there is a bleeding risk 3. Mulpleta will be used for 7 days starting 8 to 14 days prior to the procedure and discontinued 2 to 8 days prior to the procedure

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

MYALEPT

MEDICATION(S)

MYALEPT

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Pt has baseline leptin levels of less than 8 ng/mL for males OR less than 12 ng/mL for females, 2. Pt has ONE of the following: a. Diagnosis of diabetes and is being treated with Metformin AND at least one other antidiabetic agent, b. Diagnosis of hypertriglyceridemia and is being treated with at least ONE antihyperlipidemic agent, 3. Reauth: a. Pt has been screened for the presence of anti-metrelleptin antibodies, b. If presence of anti-metrelleptin antibodies, pt must still be receiving benefit from Myalept therapy, c. Pt shows improvement in hemoglobin A1c OR fasting triglyceride level

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Initial: 6 months, Reauth: Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

NARCOLEPSY

MEDICATION(S)

WAKIX

PA INDICATION INDICATOR

2 - Some FDA-Approved Indications Only

OFF LABEL USES

N/A

EXCLUSION CRITERIA

Patients with severe hepatic impairment

REQUIRED MEDICAL INFORMATION

1. For narcolepsy with cataplexy (please note Wakix is only covered for the narcolepsy with cataplexy indication) a. Patient has been diagnosed by a board certified sleep, pulmonology, or neurology specialist b. Patient exhibits symptoms of cataplexy 2. Reauth: a. For narcolepsy with cataplexy: i. Decrease in cataplexy episodes

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

Board certified in sleep, pulmonology, or neurology

COVERAGE DURATION

Plan year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

NERLYNX

MEDICATION(S)

NERLYNX

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Breast cancer dx: a. Tx of early stage HER2-positive breast cancer following adjuvant trastuzumab-based therapy OR b. Tx of HER2-positive breast cancer, in combination with capecitabine, in patients who have received 2 or more prior regimens for metastatic disease

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

NEULASTA

MEDICATION(S)

FYLNETRA, NEULASTA, NYVEPRIA, STIMUFEND, ZIEXTENZO

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of Neutropenia

a. If requesting Fylnetra, Neulasta, Nyvepria, Stimufend, or Ziextenzo, pt has trial on BOTH of the following: a. Fulphila, b. Udenyca

2. For Neulasta, Stimufend, Ziextenzo: Dx of acute hematopoietic radiation injury syndrome

a. Pt has trial of Udenyca

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

BvsD Determination

PART B PREREQUISITE

N/A

NEXAVAR

MEDICATION(S)

SORAFENIB TOSYLATE

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Used as monotherapy, 2. For HCC dx: a. Treatment for unresectable tumor or recurrent disease, 3. For Thyroid carcinoma dx: a. Tumor is refractory to treatment with iodine

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

NEXLETOL

MEDICATION(S)

NEXLETOL

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of familial hypercholesterolemia or established atherosclerotic cardiovascular disease

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

NEXLIZET

MEDICATION(S)

NEXLIZET

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of familial hypercholesterolemia or established atherosclerotic cardiovascular disease

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

NEXPLANON

MEDICATION(S)

NEXPLANON

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Use for contraception

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

NICOTROL

MEDICATION(S)

NICOTROL NS

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Pt has tried varenicline or bupropion OR has a contraindication to the use of varenicline or bupropion (established seizure disorder, concurrent anorexia or bulimia, concurrent use of MAOIs, etc)

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

NINLARO

MEDICATION(S)

NINLARO

PA INDICATION INDICATOR

4 - All FDA-Approved Indications, Some Medically-Accepted Indications

OFF LABEL USES

Autologous Stem Cell Transplant

EXCLUSION CRITERIA

1. Pt is refractory to lenalidomide or proteasome inhibitor therapy

REQUIRED MEDICAL INFORMATION

1. Multiple Myeloma A. Combination with lenalidomide (Revlimid) and dexamethasone B. Pt has previous trial on at least ONE other therapy 2. Maintenance therapy after Autologous Stem Cell Transplant

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

NITROFURAN

MEDICATION(S)

NITROFURANTOIN 25 MG/5ML SUSPENSION, NITROFURANTOIN 50 MG/10ML SUSPENSION

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Unable to swallow nitrofurantoin capsules

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

NORTHERA

MEDICATION(S)

DROXIDOPA

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Pt has previous trial on BOTH of the following: a. Midorine, b. Fludrocortisone

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

NOXAFIL

MEDICATION(S)

NOXAFIL 300 MG PACKET, POSACONAZOLE 100 MG TAB DR, POSACONAZOLE 40 MG/ML SUSPENSION

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. For treatment of invasive aspergillus 2. For aspergillus or candida prophylaxis: a. Pt is at high risk of developing infections secondary to being severely immunocompromised, 3. For oropharyngeal candidiasis: a. Pt has previous failure on BOTH of the following: 1) Itraconazole, 2) Fluconazole

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

6 Months

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

NPB INSULIN

MEDICATION(S)

INSULIN DEGLUDEC, INSULIN DEGLUDEC FLEXTOUCH

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Is the member younger than 6 years of age? (If under 6, no further questions required) 2. Patient has failed BOTH Lantus & Toujeo or has documented intolerance to Lantus and/or Toujeo?

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

NUBEQA

MEDICATION(S)

NUBEQA

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1)Diagnosis of non-metastatic castration-resistant prostate cancer (nmCRPC)

a)Patient has PSA doubling time of less than 10 months

b)Patient has tried and failed Xtandi

2)Diagnosis of metastatic hormone-sensitive prostate cancer (mHSPC)

a)Patient has tried and failed abiraterone 250mg AND Xtandi

b)Given in combination with docetaxel

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

NUPLAZID

MEDICATION(S)

NUPLAZID

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Initial: a. Onset of psychosis took place after the diagnosis of Parkinson's disease, b. Pt has previous trial on treatment with clozapine or quetiapine, 2. Reauth: a. Pt experienced a decrease in psychosis related symptoms while on treatment

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Initial: 3 months, Reauth: Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

NURTEC

MEDICATION(S)

NURTEC

PA INDICATION INDICATOR

2 - Some FDA-Approved Indications Only

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. For Acute Treatment of Migraines (Please note: Nurtec is only covered for acute treatment of migraines, prophylaxis dosing is not covered): a. Patient has tried 2 generic triptan medications OR has a contraindication to the use of triptans (e.g. established cardiovascular disease or cerebrovascular disease)

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

OCTREOTIDE

MEDICATION(S)

OCTREOTIDE ACETATE 100 MCG/ML SOLUTION, OCTREOTIDE ACETATE 1000 MCG/ML SOLUTION, OCTREOTIDE ACETATE 200 MCG/ML SOLUTION, OCTREOTIDE ACETATE 50 MCG/ML SOLUTION, OCTREOTIDE ACETATE 500 MCG/ML SOLUTION

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. For acromegaly:
 - A. Has patient failed at least TWO of the following:
 - i. Surgical resection,
 - ii. Pituitary irradiation,
 - iii. Bromocriptine,
2. Dx of metastatic carcinoid tumors:
- c. Dx of vasoactive intestinal peptide secreting tumors (VIPoma)

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan year

OTHER CRITERIA

BvsD determination

PART B PREREQUISITE

N/A

OFEV

MEDICATION(S)

OFEV

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx: a. Chronic fibrosing interstitial lung disease (ILD) with a progressive phenotype, idiopathic pulmonary fibrosis (IPF), or scleroderma (systemic sclerosis)-associated interstitial lung disease (SSc-ILD) 2. Dx has been confirmed via high-resolution computed tomography scan and/or lung biopsy, 3. Forced Vital Capacity (FVC) greater than 40 percent predicted value 4. Carbon monoxide diffusing capacity (DLCO) greater than 30 percent predicted value 5. Liver function tests (ALT, AST, and bilirubin) completed prior to initiating therapy 6. Reauth: a. A repeat liver function test has been performed after 3 months of therapy has been completed

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Initial: 3 mo, Reauth: Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

OGSIVEO

MEDICATION(S)

OGSIVEO

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of desmoid tumor/aggressive fibromatosis with documentation of tumor progression 2. Contraindication, intolerance, or failure of sorafenib

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

OJEMDA

MEDICATION(S)

OJEMDA

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

1. Patient with tumors harboring additional activating molecular alteration(s) (e.g. IDH1/2 mutations, FGFR mutations, etc) or patients with known or suspected diagnosis of neurofibromatosis type 1 (NF1)

REQUIRED MEDICAL INFORMATION

1. Dx of relapsed or refractory pediatric low-grade glioma (LGG) harboring an activating BRAF alteration based on local laboratory testing
2. At least one measurable lesion as defined by RANO 2010 criteria
3. Pt has received at least one line of prior systemic therapy and had documented evidence of radiographic progression
4. Pt has contraindication, intolerance, or failure of Tafenlar and Mekinist if BRAF V600 positive

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

OJJAARA

MEDICATION(S)

OJJAARA

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of intermediate or high-risk myelofibrosis (MF), including primary MF or secondary MF (postpolycythemia vera (PV) and post-essential thrombocythemia (ET)) 2. Hemoglobin less than 10g/dL 3. Member has tried and failed or has intolerance to Jakafi

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

ONFI

MEDICATION(S)

SYMPAZAN

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Pt has previous trail on at least TWO AED medications

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

ONUREG

MEDICATION(S)

ONUREG

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of one of: A. acute myeloid leukemia (AML) who achieved first complete remission (CR) B. AML in complete remission with incomplete blood count recovery (CRi) following intensive induction chemotherapy and the member is not able to complete intensive curative therapy

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

ORFADIN

MEDICATION(S)

NITISINONE, NITYR, ORFADIN 4 MG/ML SUSPENSION

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Initial:

- a. Diagnosis of hereditary tyrosinemia type 1 confirmed by biochemical or DNA testing
- b. Pt had a baseline succinylacetone (SA) level drawn
- c. Pt had a baseline liver function testing performed
- d. For Nityr tablets: Patient has tried and failed nitisinone capsules

2. Reauthorization:

- a. There is laboratory documentation of SA suppression on treatment when compared to baseline level

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

ORGOVYX

MEDICATION(S)

ORGOVYX

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

1. Previous treatment with another GnRH/LHR agonist/antagonist

REQUIRED MEDICAL INFORMATION

1. Dx of castrate-sensitive metastatic prostate cancer 2. Dx of metastatic disease has been confirmed by bone scan, ultrasound, CT, MRI, or biopsy 3. Serum PSA is elevated 4. Contraindication or inability to take other GNRH/LHR agonist/antagonist medication due to one of: A. Short term (6 month) use in men at risk of toxicities from standard androgen deprivation therapy (ADT) B. Intermittent ADT in frail patients at risk of ADT toxicities C. Significant underlying cardiac risk factors

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

ORIAHNN

MEDICATION(S)

MYFEMBREE, ORIAHNN

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

Maximum lifetime duration 24 months

REQUIRED MEDICAL INFORMATION

- 1)For Oriahnn and Myfembree: Heavy menstrual bleeding associated with uterine leiomyomas (fibroids)
- 2)For Myfembree: Moderate to Severe Pain Associated with Endometriosis
- 3)Premenopausal
- 4)Previous use of a combination oral contraceptive
- 5)Previous use of a progestin-only contraceptive

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

ORILISSA

MEDICATION(S)

ORILISSA

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1) previous use of combination oral contraceptive 2) previous use of progestin-only contraceptive

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year Orilissa 150 mg, maximum lifetime duration 24 months 6 months Orilissa 200 mg

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

ORKAMBI

MEDICATION(S)

ORKAMBI

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Initial A. Pt genotyped by an FDA-cleared CF mutation test B. Pt has cystic fibrosis with the homozygous F508del mutation in the CTFR gene that has been confirmed by an FDA approved test, C. Pt has FEV1 between 40-90% 2. Reauth A. Pt has been reassessed since starting therapy, B. Pt's FEV1 has increased since starting therapy

AGE RESTRICTION

Tablet: 6 years old or older, Granules: 1 year old to 5 years old

PRESCRIBER RESTRICTION

Specialist in Cystic Fibrosis or Pulmonologist

COVERAGE DURATION

Initial: 3 months Reauth: Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

ORSERDU

MEDICATION(S)

ORSERDU

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Postmenopausal female or male with a diagnosis of advanced or metastatic ER+, HER2- breast cancer 2. Confirmation of ESR1-mutated breast cancer 3. The member has experienced disease progression following at least one line of endocrine therapy

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

OSTEOPOROSIS

MEDICATION(S)

EVENITY, TERIPARATIDE (RECOMBINANT) 620 MCG/2.48ML SOLN PEN, TYMLOS

PENDING CMS APPROVAL

PAH

MEDICATION(S)

SILDENAFIL CITRATE 10 MG/ML RECON SUSP, SILDENAFIL CITRATE 20 MG TAB, TADALAFIL (PAH), TADLIQ

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Initial: a. Pt is diagnosed with pulmonary arterial hypertension, confirmed by right heart catheterization b. Pt with positive vasoreactivity test: i. Pt has contraindications to or failed maximum tolerated doses of calcium channel blockers, c. For Tadliq and sildenafil oral suspension requests: Inability to swallow tablets 2. Reauth: a. Pt has been reassessed within the past 6 months

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

PALYNZIQ

MEDICATION(S)

PALYNZIQ

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. If PKU dx: Pt has trial on sapropterin therapy 2. All dx: Pt has a blood phenylalanine (Phe) concentration of greater than 600 micromol/L

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

PANRETIN

MEDICATION(S)

PANRETIN

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

1. Systemic anti-KS therapy is required (e.g., more than 10 new KS lesions in the prior month, symptomatic lymphedema, symptomatic pulmonary KS, or symptomatic visceral involvement)

REQUIRED MEDICAL INFORMATION

1. Dx of cutaneous lesions in patients with AIDS-related Kaposi's Sarcoma. 2. Reauthorization: Patient is stable on therapy

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Initial: 6 months Reauth: Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

PART D VS PART B

MEDICATION(S)

ABILIFY ASIMTUFII, ABILIFY MAINTENA, ACETYLCYSTEINE 10 % SOLUTION, ACETYLCYSTEINE 20 % SOLUTION, ALBUTEROL SULFATE (2.5 MG/3ML) 0.083% NEBU SOLN, ALBUTEROL SULFATE 0.63 MG/3ML NEBU SOLN, ALBUTEROL SULFATE 1.25 MG/3ML NEBU SOLN, APREPITANT, ARANESP (ALBUMIN FREE), ARFORMOTEROL TARTRATE, ARISTADA, ARISTADA INITIO, ASTAGRAF XL, AZATHIOPRINE, BUDESONIDE 0.25 MG/2ML SUSPENSION, BUDESONIDE 0.5 MG/2ML SUSPENSION, BUDESONIDE 1 MG/2ML SUSPENSION, CROMOLYN SODIUM 20 MG/2ML NEBU SOLN, CYCLOPHOSPHAMIDE 25 MG CAP, CYCLOPHOSPHAMIDE 25 MG TAB, CYCLOPHOSPHAMIDE 50 MG CAP, CYCLOPHOSPHAMIDE 50 MG TAB, CYCLOSPORINE 100 MG CAP, CYCLOSPORINE 25 MG CAP, CYCLOSPORINE MODIFIED, DEPO-TESTOSTERONE, ELIGARD 22.5 MG KIT, ELIGARD 30 MG KIT, ELIGARD 7.5 MG KIT, ENGERIX-B, ENVARSUS XR, EPOGEN, EVEROLIMUS 0.25 MG TAB, EVEROLIMUS 0.5 MG TAB, EVEROLIMUS 0.75 MG TAB, EVEROLIMUS 1 MG TAB, FIRMAGON, FIRMAGON (240 MG DOSE), FLUPHENAZINE DECANOATE, FLUPHENAZINE HCL 2.5 MG/ML SOLUTION, FORMOTEROL FUMARATE, FULPHILA, GENGRAF, GRANISETRON HCL 1 MG TAB, GRANIX, HALOPERIDOL DECANOATE, HALOPERIDOL LACTATE 5 MG/ML SOLUTION, HEPLISAV-B, INVEGA HAFYERA, INVEGA SUSTENNA, INVEGA TRINZA, IPRATROPIUM BROMIDE 0.02 % SOLUTION, IPRATROPIUM-ALBUTEROL, LEUPROLIDE ACETATE 22.5 MG INJECTABLE, LEVALBUTEROL HCL, LILETTA (52 MG), LUPRON DEPOT (1-MONTH), LUPRON DEPOT (3-MONTH), LUPRON DEPOT (4-MONTH), LUPRON DEPOT (6-MONTH), LUPRON DEPOT-PED (1-MONTH) 7.5 MG KIT, LUPRON DEPOT-PED (3-MONTH) 11.25 MG (PED) KIT, METHOTREXATE SODIUM 250 MG/10ML SOLUTION, METHOTREXATE SODIUM 50 MG/2ML SOLUTION, METHOTREXATE SODIUM (PF), MICAFUNGIN SODIUM, MYCOPHENOLATE MOFETIL 200 MG/ML RECON SUSP, MYCOPHENOLATE MOFETIL 250 MG CAP, MYCOPHENOLATE MOFETIL 500 MG TAB, MYCOPHENOLATE SODIUM, MYCOPHENOLIC ACID, NIVESTYM, OLANZAPINE 10 MG RECON SOLN, ONDANSETRON 4 MG TAB DISP, ONDANSETRON 8 MG TAB DISP, ONDANSETRON HCL 4 MG TAB, ONDANSETRON HCL 4 MG/5ML SOLUTION, ONDANSETRON HCL 8 MG TAB, PENTAMIDINE ISETHIONATE, PERSERIS, PROGRAF 0.2 MG PACKET, PROGRAF 1 MG PACKET, PROLIA, PULMOZYME, RECOMBIVAX HB, RETACRIT, RISPERIDONE MICROSPHERES ER, SIROLIMUS, SODIUM CHLORIDE 0.9 % SOLUTION, STREPTOMYCIN SULFATE, TACROLIMUS 0.5 MG CAP, TACROLIMUS 1 MG CAP, TACROLIMUS 5 MG CAP, TRELSTAR MIXJECT, TWINRIX, UDENYCA, UZEDY, VARUBI (180 MG DOSE), ZIPRASIDONE MESYLATE

DETAILS

This drug may be covered under Medicare Part B or D depending on the circumstances. Information

may need to be submitted describing the use and setting of the drug to make the determination.

PCSK

MEDICATION(S)

REPATHA, REPATHA PUSHTRONEX SYSTEM, REPATHA SURECLICK

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. The patient meets one of the following:

A. The patient has tried one high-intensity statin(e.g. atorvastatin 40-80mg, rosuvastatin 20-40mg) or a maximally tolerated statin without achieving the LDL-C goal

B. The patient experienced statin-related rhabdomyolysis with documented CK elevations greater than 10x ULN

C. The patient experienced muscle-related symptoms such as myopathy or myalgia while on two separate trials of different statin therapy that both resolved upon discontinuation of statin therapy

2. AND meets ONE of the following:

A. Dx of HoFH , untreated LDL-C greater than 500mg/dL or treated LDL-C greater than 300mg/dL

i. AND cutaneous or tendon xanthoma before age 10 years, OR

ii. Elevated LDL-C levels consistent with heterozygous FH in both parents

B. Dx of HeFH or primary hyperlipidemia with fasting LDL-C of 190 mg/dL or greater on at least two separate dates at least 3 months apart

i. AND LDL-C remains greater than 100 mg/dL despite treatment on medication therapy

C. Dx of ASCVD consisting of MI, stroke, TIA, persistent intermittent claudication, coronary intervention revascularization or angina with proven ischemia

i. AND LDL-C remains greater than 70 mg/dL despite treatment on medication therapy

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

PEMAZYRE

MEDICATION(S)

PEMAZYRE

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Previously treated, unresectable locally advanced or metastatic cholangiocarcinoma A. Fibroblast growth factor receptor 2 (FGFR2) fusion or other rearrangement as detected by an FDA-approved test
2. Relapsed or refractory myeloid/lymphoid neoplasms i. Fibroblast growth factor receptor 1 (FGFR1) rearrangement as detected by an FDA-approved test

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

PHENOXYBENZAMINE

MEDICATION(S)

METYROSINE, PHENOXYBENZAMINE HCL

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Phenoxybenzamine will be used for short-term treatment of hypertension prior to surgical removal of a pheochromocytoma 2. For metyrosine, pt has failed phenoxybenzamine

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

3 mo

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

PIQRAY

MEDICATION(S)

PIQRAY (200 MG DAILY DOSE), PIQRAY (250 MG DAILY DOSE), PIQRAY (300 MG DAILY DOSE)

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative, PIK3CA-mutated, advanced or metastatic breast cancer

- a. PIK3CA-mutation positive
- b. Receiving or previous use of an endocrine-based regimen

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

POMALYST

MEDICATION(S)

POMALYST

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Multiple myeloma dx: a. Pt has tried BOTH of the following: i. Revlimid, ii. Velcade, b. Pt has demonstrated disease progression within 60 days of completion of prior therapy 2. Kaposi sarcoma dx: a. Experienced failure of highly active antiretroviral therapy (HAART) in patient with AIDS b. HIV-negative

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

PRETOMANID

MEDICATION(S)

PRETOMANID

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Diagnosis of tuberculosis 2. Used as part of an appropriate treatment regimen (e.g. bedaquiline and linezolid)

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

6 months

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

PREVYMIS

MEDICATION(S)

PREVYMIS 240 MG TAB, PREVYMIS 480 MG TAB

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

Pt has Child-Pugh class C hepatic impairment

REQUIRED MEDICAL INFORMATION

1. For: CMV prophylaxis in HSCT recipients a. Pt is post allogenic hematopoietic stem cell transplant within the last 28 days b. Pt is a CMV-seropositive recipient [R+] 2. For: CMV prophylaxis in kidney transplant recipients a. Pt is post kidney transplant within last 7 days b. Pt is high risk (Donor CMV seropositive/Recipient CMV seronegative) 3. Medication will be discontinued on or before 200 days post-transplantation

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

200 days

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

PROMACTA

MEDICATION(S)

PROMACTA

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. For ITP dx: a. Previous failure to corticosteroids, immunoglobulins, OR splenectomy, b. Initial: Evidence of bleeding OR platelet count less than 50,000/microL, c. For Reauth: Platelet count less than 400,000/microL, 2. For Hep C with Thrombocytopenia dx: a. Platelet count less than 75,000/microL, 3. For aplastic anemia dx: a. Pt has an insufficient response to immunosuppressive therapy b. In combination with immunosuppressive therapy for severe aplastic anemia

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

6 Months

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

PURIXAN

MEDICATION(S)

PURIXAN

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Used in conjunction with a combination chemotherapy treatment regimen for ALL, 2. Pt has previous failure on mercaptopurine tablets

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

PYRUKYND

MEDICATION(S)

PYRUKYND, PYRUKYND TAPER PACK

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

Homozygous R479H mutation or 2 non-missense mutations, without the presence of another missense mutation, in the PKLR gene

REQUIRED MEDICAL INFORMATION

1. Documented pyruvate kinase deficiency (PKD), presence of at least 2 mutant alleles in PKLR gene, of which at least 1 is a missense mutation 2. 6 or more transfusions in the last 12 months A. If 5 or fewer transfusions, Hb concentration less than or equal to 10.0 g/dL

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

QINLOCK

MEDICATION(S)

QINLOCK

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Diagnosis of advanced GIST 2. Prior treatment with 3 or more kinase inhibitors, including imatinib

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

QULIPTA

MEDICATION(S)

QULIPTA

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of Episodic Migraine (4-14 migraine days per month) or Chronic Migraine (greater than 14 migraine days per month) 2. Member has trial on prophylactic therapy on at least TWO of the following: A. Anti-Epileptic (ie topiramate, valproic acid) B. Beta-blocker (ie atenolol, bisoprolol, metoprolol, nadolol, nebivolol, pindolol, propranolol, timolol) C. Anti-depressant (ie amitriptyline, nortriptyline, venlafaxine) Reauthorization Criteria: 1. Patient has had a reduction in the number of migraine days per month

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Initial: Plan Year Reauthorization: Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

RCC

MEDICATION(S)

EVEROLIMUS 10 MG TAB, EVEROLIMUS 2 MG TAB SOL, EVEROLIMUS 2.5 MG TAB, EVEROLIMUS 3 MG TAB SOL, EVEROLIMUS 5 MG TAB, EVEROLIMUS 5 MG TAB SOL, EVEROLIMUS 7.5 MG TAB, TORPENZ

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

For renal angiomyolipoma, requires immediate surgery

REQUIRED MEDICAL INFORMATION

1. For RCC dx: a. Previous failure on either sunitinib or Nexavar, 2. For SEGA or TS dx: a. Patient must require therapeutic intervention and not be a candidate for surgical resection 3. Diagnosis of progressive neuroendocrine tumors of pancreatic origin (PNET) or with well-differentiated, nonfunctional neuroendocrine tumors (NET) of gastrointestinal (GI), or lung origin which are unresectable, locally advanced or metastatic 4. Diagnosed with renal angiomyolipoma with tuberous sclerosis complex with at least one angiomyolipoma greater than or equal to 3cm where there is not an immediate need for surgery 5. Hormone receptor positive HER2-negative breast cancer a. Previous use of one of letrozole or anastrozole b. Use in combination with one of exemastane, tamoxifen, or fulvestrant

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

RETEVMO

MEDICATION(S)

RETEVMO

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Diagnosis of:

- a. Metastatic RET fusion-positive non-small cell lung cancer (NSCLC)
- b. Advanced or metastatic RET-mutant medullary thyroid cancer (MTC) in patients who require systemic therapy
- c. Advanced or metastatic RET fusion-positive thyroid cancer in patients who require systemic therapy and refractory to radioactive iodine, if appropriate
- d. Locally advanced or metastatic solid tumors with a RET gene fusion, that have progressed on or following prior systemic treatment or who have no satisfactory alternative treatment options

2. Identification of a RET gene alteration using next generation sequencing (NGS), polymerase chain reaction (PCR), or fluorescence in situ hybridization (FISH)

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Initial: 6 months Reauth: Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

REVCovi

MEDICATION(S)

REVCovi

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Diagnosis of adenosine deaminase deficiency (ADA) with Severe Combined Immunodeficiency (SCID) phenotype confirmed by one of the following: A. Deficiency of adenosine deaminase (ADA) in plasma, lysed erythrocytes, fibroblasts (cultured from amniotic fluid), or chorionic villus (less than 1% of normal) B. Increase in deoxyadenosine triphosphate (dATP) levels in erythrocyte lysates compared to laboratory standard C. Decrease in ATP concentration in erythrocytes D. Molecular genetic confirmation of mutations in both alleles of the ADA1 gene E. Positive screening by T cell receptor excision circles (TRECs) 2. Not a candidate for or has failed bone marrow transplantation (BMT) 3. Platelets greater than 50,000cell/microL 4. Reauthorization A. The patient has experienced improvement in their plasma ADA activity, red blood cell dATP levels, immune function, and/or red blood cell dAXP levels

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

REVLIMID

MEDICATION(S)

LLENALIDOMIDE, REVLIMID

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. For Multiple Myeloma dx: a. Used in combination with dexamethasone, OR b. As maintenance following autologous hematopoietic stem cell transplantation (auto HSCT) 2. For MCL: a. Pt has previous trial on bortezomib AND pt has trial on at least ONE other previous therapy 3. For transfusion-dependent anemia due to myelodysplastic syndrome a. Low or imediate-1-risk myelodysplastic syndrome (MDS) associated with a deletion 5q abnormality with or without additional cytogenetic abnormalities 4. For follicular lymphoma and Marginal zone lymphoma (MZL) a. Used in combination with a rituximab product

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

REVUFORJ

MEDICATION(S)

REVUFORJ

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

1. Patients with an 11q23 partial tandem duplication
2. Active acute promyelocytic leukemia
3. Concomitant use with strong CYP3A4 inducers

REQUIRED MEDICAL INFORMATION

1. Dx of relapsed or refractory (R/R) acute leukemia with a KMT2A translocation defined as bone marrow blasts greater than or equal to 5% of reappearance of blasts in peripheral blood
2. Pt has a QTcF less than 450msec
3. For AML, patient has been treated with greater than or equal to 1 previous therapy
4. For ALL, patient has been treated with greater than or equal to 1 previous therapy, one of which is CAR-T, or documented medical rationale explaining why it would be inappropriate treatment for the patient.

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan year

OTHER CRITERIA

N/A

PART B PREREQUISITE

YES

REXULTI

MEDICATION(S)

REXULTI

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

N/A

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

REYVOW

MEDICATION(S)

REYVOW

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Patient has tried 2 generic triptan medications OR has a contraindication to the use of triptans (e.g. established cardiovascular disease or cerebrovascular disease)

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

REZDIFFRA

MEDICATION(S)

REZDIFFRA

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

1. Patient consumes more than 2 glasses (30g) of alcohol per day
2. Patients TSH is greater than 7

REQUIRED MEDICAL INFORMATION

1. Patient must have biopsy-confirmed AND FirboScan CAP AND TE-confirmed histological MASH with stage F2 or F3 fibrosis
2. Reauthorization:
 - a. Documentation shows patient has responded adequately to therapy, as demonstrated in MASH histological resolution and/or fibrosis score improvement by at least 1 stage as determined by hepatology based on liver enzymes and FibroScan CAP

AGE RESTRICTION

18 years of age or older

PRESCRIBER RESTRICTION

Hepatologist

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

REZLIDHIA

MEDICATION(S)

REZLIDHIA

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Diagnosis of relapsed or refractory AML 2. Confirmed IDH-1 mutation

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

REZUROCK

MEDICATION(S)

REZUROCK

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

Initial Authorization 1. Dx of chronic graft-versus-host disease (chronic GVHD) a. Failure of two previous lines of systemic therapy 2. For BID dosing in the setting of a PPI: a. Member must try a 30-day trial of one of the following alternative GERD treatment modalities: i. antacids ii. H2RA (ranitidine, famotidine, etc) b. Member must have not responded to daily dosing of Rezerock following at least 30 days of therapy Reauthorization 3. Documentation of stable or improved disease

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

ROZLYTREK

MEDICATION(S)

ROZLYTREK

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Diagnosis of non-small cell lung cancer A. Has reactive oxygen species 1 positive 2. Diagnosis of Neurotrophic receptor tyrosine kinase-positive solid tumors A. Tumor is metastatic or surgical resection likely to result in severe morbidity B. Progression following previous treatment or there is not an adequate alternative treatment 3. For the oral pellets: patient has inability to swallow capsules and solution made from capsules.

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

RUBRACA

MEDICATION(S)

RUBRACA

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of recurrent ovarian cancer a. Complete or partial response to platinum-based chemotherapy 2. Rubraca will be used as monotherapy 3. Dx of mCRPC that has deleterious BRCA mutation a. Previous treatment with androgen receptor-directed therapy and a taxane-based chemotherapy

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

RYDAPT

MEDICATION(S)

RYDAPT

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

1. Used for post-consolidation therapy (maintenance) for dx of AML

REQUIRED MEDICAL INFORMATION

1. For AML dx: a. Cancer is FLT3 mutation positive 2. Dx of systemic mastocytosis a. Systemic mastocytosis is identified as aggressive systemic mastocytosis (ASM), systemic mastocytosis with associated hematological neoplasm (SM-AHN), or mast cell leukemia (MCL)

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

SABRIL

MEDICATION(S)

VIGABATRIN, VIGADRONE 500 MG PACKET, VIGAFYDE, VIGPODER

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. For solution:

a. For infantile spasms:

i. Must be used as monotherapy

b. For refractory complex partial seizures

i. Pt has inability to swallow tablets

ii. Must be used as adjunctive therapy,

iii. Must have tried at least TWO of the following:

a. Rufinamide, b. Carbamazepine, c. Celontin, d. Dilantin, e. Divalproex, f. Epitol, g. Equetro, h. Ethosuximide, i. Felbamate, j. Tiagabine, k. Lamictal, l. Lamotrigine, m. Levetiracetam, n. Pregabalin, o. Primidone, p. Oxcarbazepine, q. Phenytoin, r. Topiramate, s. Valproic Acid, t. Lacosamide, u. Zonisamide

2. For tablets:

a. Must be used as adjunctive therapy,

b. Must have tried at least TWO of the following:

i. Rufinamide, ii. Carbamazepine, iii. Celontin, iv. Dilantin, v. Divalproex, vi. Epitol, vii. Equetro, viii. Ethosuximide, ix. Felbamate, x. Tiagabine, xi. Lamictal, xii. Lamotrigine, xiii. Levetiracetam, xiv. Pregabalin, xv. Primidone, xvi. Oxcarbazepine, xvii. Phenytoin, xviii. Topiramate, xix. Valproic Acid, xx. Lacosamide, xxi. Zonisamide

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

1. Periodic vision testing

PART B PREREQUISITE

N/A

SCEMBLIX

MEDICATION(S)

SCEMBLIX

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Newly diagnosed Philadelphia chromosome-positive chronic myeloid leukemia (Ph+ CML) in chronic phase (CP)
2. Philadelphia chromosome-positive chronic myeloid leukemia (Ph+ CML) in chronic phase (CP):
 - A. Trial and failure of at least 1 TKI
3. Philadelphia chromosome-positive chronic myeloid leukemia (Ph+ CML) in chronic phase (CP) with T315I mutation:
 - A. Documentation of testing for mutation
 - B. Must try and fail ponatinib

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

SIGNIFOR

MEDICATION(S)

SIGNIFOR

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. For Cushing's:

- a. Pt is NOT a candidate for pituitary surgery,
- b. If pt previously had pituitary surgery: Pt continues to have high 24-hour urinary free cortisol levels
- c. Patient has tried at least one of the following:
 - i. ketoconazole
 - ii. metyrapone

2. For acromegaly

- a. Pt has had surgical resection of the pituitary gland OR is not a candidate for surgery/radiation therapy
- b. Pt has tried at least one of the following:
 - i. bromocriptine
 - ii. cabergoline
 - iii. octreotide acetate
- c. Pt has tried sandostatin LAR, somatuline depot, or lanreotide

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Initial: Plan year Reauth: Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

SIRTURO

MEDICATION(S)

SIRTURO

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Pt has previous failure on at least TWO of the following: a. Ethambutol, b. Streptomycin, c. Pyrazinamide, d. Amikacin/kanamycin, e. Cycloserine/terizidone, f. Ethionamide, g. Capreomycin, h. Levofloxacin, i. Moxifloxacin, j. Ofloxacin

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

6 Months

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

SOHONOS

MEDICATION(S)

SOHONOS

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of fibrodysplasia ossificans progressiva (FOP) confirmed by genetic testing (documentation must be submitted)

AGE RESTRICTION

Females age 8 years or older and Males age 10 years or older

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

SOMAVERT

MEDICATION(S)

SOMAVERT

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Pt has had surgical resection of the pituitary gland OR is not a candidate for surgery/radiation therapy, 2. Patient has tried at least ONE of the following: a. Bromocriptine, b. Cabergoline, c. Octreotide acetate 3. Patient has tried sandostatin LAR, somatuline depot, or lanreotide

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

SPRYCEL

MEDICATION(S)

DASATINIB

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Diagnosis of CML or ALL

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

STELARA

MEDICATION(S)

STELARA 45 MG/0.5ML SOLN PRSYR, STELARA 45 MG/0.5ML SOLUTION, STELARA 90 MG/ML SOLN PRSYR

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

Combination therapy with another biologc medication, JAK inhibitor, or Otezla

REQUIRED MEDICAL INFORMATION

1. Crohn's disease dx 2. Ulcerative Colitis dx 3. Plaque Psoriasis dx: Pt has previous failure with at least ONE of the following: a. methotrexate, b. cyclosporine, c. acitretin 4. Psoriatic arthritis dx: Pt has previous failure with at least ONE of the following: a. methotrexate, b. leflunomide, c. hydroxychloroquine, d. sulfasalazine, e. injectable gold, f. oral gold, g. azathioprine, h. D-penicillamine, i. cyclosporine

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

STIVARGA

MEDICATION(S)

STIVARGA

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx metastatic colorectal cancer, 2. Dx gastrointestinal stromal tumor, 3. Dx hepatocellular carcinoma

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

SUCRAID

MEDICATION(S)

SUCRAID

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx is confirmed by a Small Bowel Biopsy Disaccharidase Measurement demonstrating 2 SD or more below mean for sucrase activity with or without isomaltase activity

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

SUTENT

MEDICATION(S)

SUNITINIB MALATE

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

1. Sutent used as combination therapy with other chemotherapies

REQUIRED MEDICAL INFORMATION

1. For GIST dx: a. Disease progression or intolerance to imatinib, 2. For pNET dx: a. Tumor is unresectable locally advanced or metastatic

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

SYNAREL

MEDICATION(S)

SYNAREL

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

Pt has previous trial on leuprolide acetate

AGE RESTRICTION

1. For CPP: Treatment initiated at or before 8 years of age in girls and 9 years of age in boys 2. For Endometriosis: 18 years old or older

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

SYPRINE

MEDICATION(S)

TRIENTINE HCL

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Pt has failure on penicillamine,

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

TABRECTA

MEDICATION(S)

TABRECTA

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of metastatic non-small cell lung cancer (NSCLC) 2. Tumor mutation leading to mesenchymal-epithelial transition (MET) exon 14 skipping

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

TADALAFIL

MEDICATION(S)

TADALAFIL 5 MG TAB

PA INDICATION INDICATOR

2 - Some FDA-Approved Indications Only

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of Benign Prostatic Hypertrophy (BPH)

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

TAFINLAR

MEDICATION(S)

TAFINLAR

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1)For unresectable or metastatic melanoma dx:

a)Pt is BRAF (V600E or V600K) mutation positive

2)For adjuvant treatment of melanoma dx:

a)Pt is BRAF (V600E or V600K) mutation positive

b)Pt has failed Opdivo AND Keytruda

3)For metastatic NSCLC dx:

a)Pt is BRAF V600E positive

4)For Locally Advanced or Metastatic Anaplastic Thyroid Cancer dx:

a)Pt is BRAF V600E positive with no satisfactory locoregional treatment options

5)For Unresectable or Metastatic Solid Tumors dx:

a)Pt is BRAF V600E positive and has progressed following prior treatment and have no satisfactory alternative treatment options

6)For Low-Grade Glioma dx”

a)Pt is BRAF V600E mutation positive and requires systemic therapy

b)Pt has failed Zelboraf (with or without Cotellic)

7)For Tafinlar tablets for oral suspension: Member has inability to swallow capsules

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

YES

TAGRISSE

MEDICATION(S)

TAGRISSE

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Diagnosis of metastatic non-small cell lung cancer (NSCLC) and will be used as one of the following:

A. adjuvant therapy after tumor resection, whose tumors have epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 L858R mutations

B. locally advanced, unresectable (stage III) NSCLC whose disease has not progressed during or following concurrent or sequential platinum-based chemoradiation therapy and whose tumors have EGFR exon 19 deletions or exon 21 L858R mutations

C. first-line treatment of metastatic NSCLC whose tumors have EGFR exon 19 deletions or exon 21 L858R mutations

D. first-line treatment of locally advanced or metastatic NSCLC whose tumors have EGFR exon 19 deletions or exon 21 L858R mutations in combination with pemetrexed and platinum-based chemotherapy

E. treatment of adult patients with metastatic EGFR T790M mutation positive NSCLC, whose disease has progressed on or after EGFR TKI therapy

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

TALTZ

MEDICATION(S)

TALTZ 80 MG/ML SOLN A-INJ, TALTZ 80 MG/ML SOLN PRSYR

PA INDICATION INDICATOR

2 - Some FDA-Approved Indications Only

OFF LABEL USES

N/A

EXCLUSION CRITERIA

Combination therapy with another biologic medication, JAK inhibitor, or Otezla

REQUIRED MEDICAL INFORMATION

1. Dx of Ankylosing Spondylitis (Please note, Taltz is not covered for indications other than Ankylosing spondylitis)
2. Member has tried adalimumab and Renflexis

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

YES

TALZENNA

MEDICATION(S)

TALZENNA

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of HRR Gene-mutated mCRPC, in combination with enzalutamide (Xtandi) 2. Deleterious or suspected deleterious germline BRCA, HER2-negative locally advanced or metastatic breast cancer

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

TARCEVA

MEDICATION(S)

ERLOTINIB HCL

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. For NSCLC dx: a. Pt with EGFR mutation, b. Erlotinib is not used in combination with platinum-based chemotherapy, 2. For pancreatic cancer dx: a. Combination with gemcitabine

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

TARGRETIN

MEDICATION(S)

BEXAROTENE

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

1. If female: Pt planning to become pregnant

REQUIRED MEDICAL INFORMATION

1. Capsules: a. Pt has previous failure on at least ONE of the following: i. Antineoplastic chemotherapy, ii. Interferon alfa and gamma, iii. Interleuking-12, iv. Interleukin-2, 2. Gel: a. Pt has previous failure on at least ONE of the following: i. PUVA, ii. UVB, iii. EVT, iv. Photophoresis, v. Systemic cytotoxic chemotherapy, vi. Topical nitrogen mustard, vii. Topical carmustine

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

TARPEYO

MEDICATION(S)

FILSPARI, TARPEYO

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

Currently receiving dialysis or has undergone kidney transplant

REQUIRED MEDICAL INFORMATION

1. Dx of biopsy verified primary immunoglobulin A (IgA) nephropathy
2. Proteinuria defined as a urine protein-to-creatinine ratio (UPCR) greater than or equal to 1.5 g/g
3. Failure of at least a 3-month trial on SGLT2
4. For Tarpeyo: Failure of three months of Filspari

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

For Tarpeyo: 9 months, no reauthorization

For Filspari: Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

TASIGNA

MEDICATION(S)

TASIGNA

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

For Ph+ CML Dx: Pts who have BCR-ABL1 mutations T315I, Y253H, E255K/V, F359V/C/I, or G250E

REQUIRED MEDICAL INFORMATION

1. Adult or pediatric with newly diagnosed Philadelphia chromosome positive (Ph+) CML in chronic phase
2. Adult with chronic phase and accelerated phase Ph+ CML a. Resistant or intolerant to imatinib
3. Pediatric patient with chronic phase or accelerated phase Ph+ CML a. Resistant or intolerant to prior TKI therapy

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

TAVALISSE

MEDICATION(S)

TAVALISSE

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Initial: a. Pt has previous trial on previous therapy of at least ONE of the following: i. Corticosteroids ii. Immunoglobulins iii. Splenectomy iv. Thrombopoietin Receptor Agonist b. Platelet count is less than 50000/micL 2. Reauth: a. Platelet count is greater than 50000/micL

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Initial: 3 Months Reauth: Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

TAVNEOS

MEDICATION(S)

TAVNEOS

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

1. Patient currently requires dialysis

REQUIRED MEDICAL INFORMATION

Initial: 1. Diagnosis of ANCA-associated vasculitis 2. ANCA-antibody titer test 3. BVAS score (1 major item, 3 non-major items, or 2 renal items of proteinuria and hematuria) 4. Patient is currently receiving rituximab or cyclophosphamide Reauth: 1. Patient has experienced improvement in BVAS score

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Initial: 6 months Reauth: Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

TAZVERIK

MEDICATION(S)

TAZVERIK

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx: Metastatic or locally advanced epithelioid sarcoma a. Diagnosis confirmed by pathology b. SMARCB1/INI1 deficient tumor c. Tumor is not eligible for complete resection 2. Dx: Follicular lymphoma a. Relapsed or refractory tumor i. positive for EZH2 mutation ii. Previous use of at least 2 prior systemic therapies b. No satisfactory alternative treatment options available

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

TEPMETKO

MEDICATION(S)

TEPMETKO

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Diagnosis of metastatic non-small cell lung cancer (NSCLC) harboring mesenchymal-epithelial transition (MET) exon 14 skipping alterations.

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

TIBSOVO

MEDICATION(S)

TIBSOVO

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. AML newly diagnosed is susceptible isocitrate dehydrogenase-1 (IDH1) mutation
2. AML is susceptible isocitrate dehydrogenase-1 (IDH1) mutation and has relapsed or is refractory to one or more prior anticancer regimens
3. Locally advanced or metastatic cholangiocarcinoma with an isocitrate dehydrogenase-1 (IDH-1) mutation as detected by an FDA-approved test and is refractory to previous treatment
4. Relapsed or refractory myelodysplastic syndromes (MDS) with a susceptible isocitrate dehydrogenase-1 (IDH1) mutation

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

TOLCAPONE

MEDICATION(S)

TOLCAPONE

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. The member is currently receiving carbidopa/levodopa therapy
2. Member is experiencing `off episodes that requires adjunctive therapy and has failed at least two of the following: Entacapone, Ongentys, Dopamine agonist (pramipexole, ropinirole, etc), MAO B inhibitor (rasagiline, selegiline)

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

TRUQAP

MEDICATION(S)

TRUQAP 160 MG TAB, TRUQAP 200 MG TAB

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

1. diabetes mellitus type 1, type 2, requiring insulin treatment or HbA1c greater than or equal to 8%

REQUIRED MEDICAL INFORMATION

1. Dx of HR-positive, HER2-negative, locally advanced or metastatic breast cancer
2. Patient has PIK3CA/AKT1/PTEN-alterations.
 - A. If PIK3CA, patient must have failed Piqray
3. Patient has failed at least one endocrine-based regimen in the metastatic setting or recurrence on or within 12 months of completing adjuvant therapy
4. Truqap will be used in combination with fulvestrant

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

TUKYSA

MEDICATION(S)

TUKYSA

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. For RAS wild-type unresectable or metastatic HER2-positive breast cancer
 - a. Used in combination with trastuzumab and capecitabine
2. For unresectable or metastatic colorectal cancer
 - a. Used in combination with trastuzumab
 - b. Following treatment with fluoropyrimidine, oxaliplatin, and irinotecan based chemotherapy

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

TURALIO

MEDICATION(S)

TURALIO 125 MG CAP

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

- A. Diagnosis of Tenosynovial Giant Cell Tumor
- B. Condition is associated with severe morbidity or functional limitations
- C. Surgery will NOT improve status

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

TYENNE

MEDICATION(S)

TYENNE 162 MG/0.9ML SOLN A-INJ, TYENNE 162 MG/0.9ML SOLN PRSYR

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

1. Combination use with another biologic medications, janus kinase inhibitor (JAK), or Otezla

REQUIRED MEDICAL INFORMATION

1. Dx of:

- A. Cytokine release syndrome (CRS)
- B. Giant cell arteritis (GCA)
- C. Systemic juvenile idiopathic arthritis (JIA)
- D. Moderate to severe rheumatoid arthritis (RA)
- E. Systemic sclerosis-associated interstitial lung disease (SSc-ILD)
- F. Covid-19

2, For RA or JIA

A. Pt has failed at least three months therapy on at least ONE of the following:

i. methotrexate, ii. leflunomide, iii. hydroxychloroquine, iv. sulfasalazine, OR contraindication to use (Clinical diagnosis of alcohol use disorder, alcoholic liver disease or other chronic liver disease, Pregnancy, breastfeeding, or currently planning pregnancy, Blood dyscrasias (e.g., thrombocytopenia, leukopenia, significant anemia), Elevated liver transaminases, Hypersensitivity, or history of intolerance or adverse event, Interstitial pneumonitis or clinically significant pulmonary fibrosis, Myelodysplasia, Renal impairment

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

TYKERB

MEDICATION(S)

LAPATINIB DITOSYLATE

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. For advanced or metastatic HER-2 positive breast cancer dx:
 - a. Previous failure on anthracycline, taxane, and trastuzumab AND
 - b. Combination therapy with capecitabine
2. For postmenopausal HER-2 receptor hormone receptor positive breast cancer dx:
 - a. Combination therapy with aromatase inhibitor

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

UBRELVY

MEDICATION(S)

UBRELVY

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Patient has tried 2 generic triptan medications OR has a contraindication to the use of triptans (e.g. established cardiovascular disease or cerebrovascular disease)

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

VALCHLOR

MEDICATION(S)

VALCHLOR

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Pt has previous trial on at least ONE previous skin directed therapy of the following:
 - a. Topical corticosteroid, b. Topical carmustine, c. Topical retinoid, d. Radiation therapy, e. Phototherapy

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

VANFLYTA

MEDICATION(S)

VANFLYTA

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Newly diagnosed acute myeloid leukemia that is FLT3 internal tandem duplication (ITD)-positive as detected by an FDA approved test.
2. Will be taken in combination with standard cytarabine and anthracycline induction, cytarabine consolidation, and as maintenance monotherapy following consolidation chemotherapy

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

VAXCHORA

MEDICATION(S)

VAXCHORA

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Member is traveling to area of active cholera transmission (Please note, the dates and location of active cholera transmission area member is going to need to be documented in request)
2. Member is between the ages of 2 and 64

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

VELTASSA

MEDICATION(S)

VELTASSA

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Initial:

a. If patient is on a ACE or ARB must meet BOTH of the following:

i. Pt has been tried on a loop or thiazide diuretic or has a contraindication to one of these diuretics,

ii. The dose of the ACE or ARB has been reduced in an attempt to lower serum potassium levels,

b. Serum potassium levels above 5.1 mmol/L on two separate screenings,

c. Pt has chronic kidney disease with an eGFR of 15 to 60 mL/min

d. If patient is 18 years or older: Pt has failed Lokelma

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Initial: Plan Year

Reauth: Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

VEMLIDY

MEDICATION(S)

VEMLIDY

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

Pt has decompensated hepatic impairment

REQUIRED MEDICAL INFORMATION

Pt has previous trial on entecavir

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

VENCLEXTA

MEDICATION(S)

VENCLEXTA, VENCLEXTA STARTING PACK

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. For CLL/SLL dx
2. For AML dx: a. Pt is ineligible for induction therapy OR b. Pt is 75 years or older

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

VERQUVO

MEDICATION(S)

VERQUVO

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

1. Pregnancy

REQUIRED MEDICAL INFORMATION

1. Dx of chronic heart failure

A. New York Heart Association class II-IV

B. Left ventricular ejection fraction less than 45%

2. Previous hospitalization due to heart failure within the last 6 months or outpatient IV diuretic treatment within the last 3 months

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

VERZENIO

MEDICATION(S)

VERZENIO

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Adjuvant treatment of HR-positive, HER-2 negative, node positive early breast cancer at high risk of recurrence in combination with tamoxifen or aromatase inhibitor
2. Dx of HR-positive, HER2-negative Advanced or Metastatic breast cancer: Must meet a, b, OR c of the following:
 - a. Pt is receiving Verzenio in combination with an aromatase inhibitor as initial endocrine-based therapy
 - b. Pt with disease progression following endocrine therapy in combination with fulvestrant,
 - c. Pt with disease progression following endocrine therapy and prior chemotherapy in the metastatic setting

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

VIJOICE

MEDICATION(S)

VIJOICE

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Confirmed/documented diagnosis of PIK3CA Related Overgrowth Spectrum (PROS)
 - a. Patient has mutation in the PIK3CA gene.

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

VITRAKVI

MEDICATION(S)

VITRAKVI

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. solid tumor with a NTRK gene fusion
2. Metastatic or unable to have surgery
3. Received previous treatment

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

VIVJOA

MEDICATION(S)

VIVJOA

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Hx of recurrent vulvovaginal candidiasis, defined as at least 3 acute episodes in the last 12 months
2. Patient must be one of the following:
 - A. Post-menopausal
 - B. Not of reproductive potential (i.e. tubal ligation, hysterectomy, etc)
3. Patient has experienced a recurrence during or following 6 months of oral fluconazole maintenance therapy

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

14 weeks

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

VIZIMPRO

MEDICATION(S)

VIZIMPRO

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1) First line therapy 2) EGFR exon 19 deletion or EFGR exon 21 L858R substitution

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

VONJO

MEDICATION(S)

VONJO

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Diagnosis of intermediate or high-risk primary or secondary myelofibrosis (MF)
 - a. For secondary MF: post-polycythemia vera or post-essential thrombocythemia
2. Documentation showing platelet counts below 50,000/mm³ within the last 30 days
3. For intermediate risk: Inadequate response or intolerance to hydroxyurea, Pegasys, or Jakafi
4. For high risk: patient is not a candidate for transplant
5. Reauthorization: CBC and platelet count required. If above 50,000mm³, Vonjo is no longer indicated. Jakafi is approved for use in patients with platelet counts above 50,000/mm³

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Initial: 6 months

Reauthorization: Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

VORANIGO

MEDICATION(S)

VORANIGO

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Initial:

- A. Patient has Grade 2 oligodendroglioma or astrocytoma per WHO criteria
- B. Patient has confirmed IDH1 or IDH2 gene mutation
- C. Patient has had at least 1 prior surgery for glioma (biopsy, sub-total resection, gross-total resection)
- D. One of:
 - i. Patient has not received other prior anticancer therapy, including chemotherapy and radiotherapy
 - ii. Patient has received both RT and chemotherapy AND
 - a. patient has KPS greater than or equal to 60 AND
 - b. For IDH1 confirmed mutations in adult patients, patient has failed Tibsovo
- E. Patient has MRI evaluable, measurable, non-enhancing disease

2. Reauthorization

- A. Patient does not have imaging-based (MRI) disease progression

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

VOSEVI

MEDICATION(S)

VOSEVI

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Confirmation of genotype

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

12 Weeks

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

VOTRIENT

MEDICATION(S)

PAZOPANIB HCL

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Soft Tissue Sarcoma dx:
 - a. Previous trial on at least ONE prior therapy
2. Advanced renal cell carcinoma dx

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

VOXZOGO

MEDICATION(S)

VOXZOGO

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of achondroplasia, confirmed by genetic testing
2. Ambulatory and able to stand without assistance
3. Member has open epiphyses
4. Reauthorization
 - a) No evidence of growth plate closure (proximal tibia, distal femur)

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

VYNDAQEL

MEDICATION(S)

VYNDAMAX, VYNDAQEL

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

1. Use in combination with Onpattro or Tegsedi

REQUIRED MEDICAL INFORMATION

I. For cardiomyopathy of hereditary transthyretin-mediated amyloidosis (ATTR-CM)

A. Applicable mutation to the transthyretin (TTR) gene (e.g., Val122Ile, Thr60Ala, or Ile68Leu)

B. Clinical manifestations of amyloid fibrosis in the myocardium (e.g., AV bundle branch block, restrictive cardiomyopathy and heart failure with preserved ejection fraction [HFpEF])

II. For cardiomyopathy of wild type transthyretin-mediated amyloidosis (ATTR-CM)

B. Negative for monoclonal protein in blood or urine (evaluated by serum free light chains and serum protein electrophoresis with urine immunofixation)

C. Absence of monoclonal proteins has the patient undergone nuclear scintigraphy diagnosis OR presence of monoclonal proteins has the patient undergone tissue biopsy to rule out light chain cardiac amyloidosis

D. Clinical manifestations of amyloid fibrosis in the myocardium (e.g., AV bundle branch block, restrictive cardiomyopathy and heart failure with preserved ejection fraction [HFpEF])

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

WEGOVY

MEDICATION(S)

WEGOVY

PA INDICATION INDICATOR

2 - Some FDA-Approved Indications Only

OFF LABEL USES

N/A

EXCLUSION CRITERIA

1. NYHA Class IV Heart Failure
2. Dx of type 1 or type 2 diabetes
3. Combination use with another GLP-1 Receptor Agonist (tirzepatide, dulaglutide, etc)

REQUIRED MEDICAL INFORMATION

1. Established cardiovascular disease defined as history of one of the following:
 - a. Prior myocardial infarction
 - b. Prior stroke
 - c. Symptomatic peripheral arterial disease as evidenced by intermittent claudication with ankle-brachial index greater than 0.85, peripheral arterial revascularization procedure, or amputation due to atherosclerotic disease
2. BMI of 27 or greater

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

Cardiologist

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

WELIREG

MEDICATION(S)

WELIREG

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

1.Requires immediate surgery

REQUIRED MEDICAL INFORMATION

1.Dx von Hippel-Lindau (VHL) disease who require therapy for associated:

- a. renal cell carcinoma (RCC)
- b. central nervous system (CNS) hemangioblastomas
- c. pancreatic neuroendocrine tumors (pNET)

2. Dx of advanced renal cell carcinoma (RCC)

a. Will be used following a programmed death receptor-1 (PD-1) or programmed death-ligand 1 (PD-L1) inhibitor and a vascular endothelial growth factor tyrosine kinase inhibitor (VEGF-TKI)

3. Reauthorization: Patient is stable on therapy

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

WINREVAIR

MEDICATION(S)

WINREVAIR

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

1. Diagnosis of pulmonary arterial hypertension WHO Groups 2, 3, 4, or 5
2. Diagnosis of the following PAH Group 1 subtypes: human immunodeficiency virus (HIV)-associated PAH and PAH associated with portal hypertension. Exclusions in PAH group 1 should also include schistosomiasis associated PAH and pulmonary veno-occlusive disease

REQUIRED MEDICAL INFORMATION

1. Dx of pulmonary arterial hypertension (PAH), classified as WHO functional class (FC) II or III
2. Documented diagnostic right heart catheterization confirming the diagnosis of World Health Organization (WHO) pulmonary arterial hypertension (PAH) Group1 in any of the following subtypes:
 - a. Idiopathic PAH
 - b. Heritable PAH
 - c. Drug/Toxin-induced PAH
 - d. PAH associated with connective tissue disorder
 - e. PAH associated with simple, congenital tissue disease
3. Patient has undergone vasoreactivity testing and failed maximum tolerated doses of calcium channel blockers (if applicable)
4. Pulmonary capillary wedge pressure (PCWP) or left ventricular end-diastolic pressure of less than or equal to 15 mmHg
5. Patients current therapy consists of 2 background PAH medications including:
 - a. PDE-5 inhibitor (tadalafil, sildenafil), plus endothelin receptor antagonist (ambrisentan, bosentan, Opsumit), and a prostacyclin agonist (treprostinil or epoprostenol, or selexipag)
6. Patient has a documented intolerance or disease progression on a three-drug regimen

AGE RESTRICTION

18 years of age or older

PRESCRIBER RESTRICTION

Pulmonologist or Cardiologist

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

XALKORI

MEDICATION(S)

XALKORI

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1) Dx with:

- a) metastatic non-small cell lung cancer (NSCLC) whose tumors are anaplastic lymphoma kinase (ALK)
- b) metastatic non-small cell lung cancer (NSCLC) whose tumors are ROS1-positive
- c) relapsed or refractory, systemic anaplastic large cell lymphoma (ALCL) that is ALK-positive
- d) unresectable, recurrent, or refractory ALK-positive inflammatory myofibroblastic tumor

2) For NSCLC and ALK-positive inflammatory myofibroblastic tumor: Patient has failed/intolerance/contraindication to Alecensa

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

XDEMZY

MEDICATION(S)

XDEMZY

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of Demodex blepharitis

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

Ophthalmologist

COVERAGE DURATION

6 weeks

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

XELJANZ

MEDICATION(S)

XELJANZ, XELJANZ XR

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

1. Combination therapy with another biologic medication, JAK inhibitor, or Otezla

REQUIRED MEDICAL INFORMATION

1. Dx of RA, JIA, PsA: Pt has failed at least three months therapy on at least ONE of the following:
 - a. methotrexate, b. leflunomide, c. hydroxychloroquine, d. sulfasalazine, OR contraindication to use (Clinical diagnosis of alcohol use disorder, alcoholic liver disease or other chronic liver disease, Pregnancy, breastfeeding, or currently planning pregnancy, Blood dyscrasias (e.g., thrombocytopenia, leukopenia, significant anemia), Elevated liver transaminases, Hypersensitivity, or history of intolerance or adverse event, Interstitial pneumonitis or clinically significant pulmonary fibrosis, Myelodysplasia, Renal impairment
2. Dx of RA, JIA, PsA, UC, AS: Pt has failed a TNF inhibitor

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

XENAZINE

MEDICATION(S)

TETRABENAZINE

PA INDICATION INDICATOR

4 - All FDA-Approved Indications, Some Medically-Accepted Indications

OFF LABEL USES

Tardive Dyskinesia

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Diagnosis of chorea due to Huntington's Disease OR
2. Diagnosis of Tardive Dyskinesia

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

XERMELO

MEDICATION(S)

XERMELO

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Initial:

- a. Pt has more than 4 bowel movements a day despite treatment with sandostatin analog therapy for at least 3 months
- b. Pt has previous trial of lomotil AND loperamide
- c. Xermelo will be used in combination with sandostatin analog

2. Reauth:

- a. Pt has experienced improvement in bowel movement frequency since starting Xermelo

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

6 Months

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

XGEVA

MEDICATION(S)

XGEVA

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Must meet at least ONE of the following:

- a. Pt has giant cell tumor of the bone that is unresectable or where surgical resection is likely to result in severe morbidity,
- b. Pt has a diagnosis of bone metastases related to a solid tumor,
- c. Pt has a diagnosis of metastatic breast or prostate cancer,
- d. Pt has previously been treated with Zometa and had disease progression OR adverse reaction to the treatment

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

BvsD determination

PART B PREREQUISITE

YES

XHANCE

MEDICATION(S)

XHANCE

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Diagnosis of Chronic Rhinosinusitis with or without Nasal Polyps.
2. Previous use of mometasone intranasal spray

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

XIFAXAN

MEDICATION(S)

XIFAXAN

PA INDICATION INDICATOR

4 - All FDA-Approved Indications, Some Medically-Accepted Indications

OFF LABEL USES

1. Clostridium difficile infection

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Diagnosis of:

- a. Clostridium difficile (C. diff.) infection
- b. Hepatic encephalopathy (HE)
- c. Irritable bowel syndrome (IBS)
- d. Traveler's diarrhea

2. Diagnosis of Hepatic encephalopathy (HE)

- a. Previous failure on or has intolerance to lactulose therapy

3. Diagnosis of Irritable bowel syndrome (IBS)

- a. Previous failure of at least TWO antispasmodic or antibiotic treatments (e.g., amoxicillin-clavulanate, cephalexin, ciprofloxacin, dicyclomine, doxycycline, gentamicin, metronidazole, neomycin, trimethoprim-sulfamethoxazole)

4. Diagnosis of Clostridium difficile (C. diff) infection

- a. Patient has experienced relapse after prior use of oral vancomycin

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

XOLAIR

MEDICATION(S)

XOLAIR

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

1. Concurrent therapy with another biologic medication
2. Combination use with Palforzia

REQUIRED MEDICAL INFORMATION

1. For allergic asthma dx:
 - a. Patient has tried at least 3 months on any of the following combinations:
 - i. ICS/LABA
 - ii. ICS/LTRA
 - iii. ICS/LAMA
 - iv. ICS/LABA/LAMA (Trelegy)
 - b. Pt has failed had two or more exacerbations requiring oral corticosteroids OR one exacerbation that led to a hospitalization within the past 12 months
 - c. Pt's IgE level is greater than or equal to 30 IU/mL
 - d. Pt is less than 330 lbs. (150 kg),
2. For Chronic Spontaneous Urticaria (CSU) dx:
 - a. Pt has previous failure on a H-1 antagonist, (e.g., cetirizine, hydroxyzine)
 - b. Pt has previous failure of leukotriene receptor antagonist (LTRA) (e.g., montelukast)
3. Dx of Nasal Polyps
 - a. Previous use of systemic corticosteroid therapy
 - b. Patient has had previous failure of and will be used in combination with nasal corticosteroids
4. IgE-mediated Food Allergy
 - a. Documentation supports that the patient has a history of Type 1 (IgE-mediated) food allergy
 - b. Diagnosis is confirmed through positive skin prick test or positive serum IgE
 - c. Body weight and documented pretreatment IgE levels are within the dosing recommendations based on the FDA prescribed information

- d. Patient will continue with food allergen avoidance where appropriate
- e. For reauthorization:
 - i. Documentation must show the patient has experienced a significant reduction in frequency or severity of incidental exposure reactions

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

For Food Allergy: Allergist or Immunologist

COVERAGE DURATION

For Food Allergy: 6 months, Reauth: Plan Year

For other indications: Plan Year

OTHER CRITERIA

BvsD Determination

PART B PREREQUISITE

N/A

XOSPATA

MEDICATION(S)

XOSPATA

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Relapsed or refractory AML
2. patient has a FLT3 mutation detected by an FDA-approved test

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

XPOVIO

MEDICATION(S)

XPOVIO (100 MG ONCE WEEKLY) 50 MG TAB THPK, XPOVIO (40 MG ONCE WEEKLY) 40 MG TAB THPK, XPOVIO (40 MG TWICE WEEKLY) 40 MG TAB THPK, XPOVIO (60 MG ONCE WEEKLY) 60 MG TAB THPK, XPOVIO (60 MG TWICE WEEKLY), XPOVIO (80 MG ONCE WEEKLY) 40 MG TAB THPK, XPOVIO (80 MG TWICE WEEKLY)

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of relapsed or refractory multiple myeloma
 - a. Previous use of at least one prior therapies
2. Dx of diffuse large B-cell lymphoma
 - a. Previous use of at least two lines of systemic therapy

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

XTANDI

MEDICATION(S)

XTANDI

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

- 1) Diagnosis of non-metastatic castration sensitive prostate cancer (nmCSPC)
 - a) Patient has failed androgen deprivation therapy (ADT) with one of the following: orchiectomy, LHRH agonist (leuprolide, Lupron, Zoladex), LHRH antagonist (Firmagon)
- 2) Diagnosis of metastatic castration-sensitive prostate cancer (mCSPC)
 - a) Patient has failed ADT and abiraterone 250mg
- 3) Diagnosis of non-metastatic castration-resistant prostate cancer (nmCRPC)
 - a) Patient has PSA doubling time of less than 10 months

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

YORVIPATH

MEDICATION(S)

YORVIPATH

PENDING CMS APPROVAL

ZARXIO

MEDICATION(S)

NEUPOGEN, RELEUKO 300 MCG/0.5ML SOLN PRSYR, RELEUKO 480 MCG/0.8ML SOLN PRSYR, ZARXIO

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1)Dx of Neutropenia

2)If requesting Neupogen, Releuko or Zarxio, pt has trial on BOTH of the following: a. Nivestym, b. Granix

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

ZEJULA

MEDICATION(S)

ZEJULA 100 MG TAB, ZEJULA 200 MG TAB, ZEJULA 300 MG TAB

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Advanced epithelial ovarian, fallopian tube, or primary peritoneal cancer
 - a. Complete or partial response to first-line platinum-based chemotherapy
2. Deleterious or suspected deleterious germline BRCA-mutated (gBRCAmut) recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer
 - a. Complete or partial response to platinum-based chemotherapy

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

ZELAPAR

MEDICATION(S)

ZELAPAR

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Pt diagnosed with Parkinson's
2. Pt is treated with carbidopa/levodopa and will continue to be treated with carbidopa/levodopa in combination with Zelapar
3. Pt has failed generic selegiline

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

ZELBORAF

MEDICATION(S)

COTELLIC, ZELBORAF

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. For Metastatic Melanoma dx:
 - a. Pt is BRAF V600E positive for Zelboraf monotherapy OR
 - b. BRAF V600E or V600K positive for Zelboraf plus Cotellic
2. For Erdheim-Chester Disease:
 - a. Zelboraf monotherapy
 - b. Pt is BRAF V600 positive

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

ZEPBOUND

MEDICATION(S)

ZEPBOUND 10 MG/0.5ML SOLN A-INJ, ZEPBOUND 12.5 MG/0.5ML SOLN A-INJ, ZEPBOUND 15 MG/0.5ML SOLN A-INJ, ZEPBOUND 2.5 MG/0.5ML SOLN A-INJ, ZEPBOUND 5 MG/0.5ML SOLN A-INJ, ZEPBOUND 7.5 MG/0.5ML SOLN A-INJ

PENDING CMS APPROVAL

ZOLINZA

MEDICATION(S)

ZOLINZA

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Pt has progressive, persistent, or recurrent disease,
2. Pt has tried at least TWO prior systemic therapies

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

ZONISADE

MEDICATION(S)

ZONISADE

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of partial-onset seizures
2. Inability to swallow tablets and capsules

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

ZTALMY

MEDICATION(S)

ZTALMY

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Confirmation of CDKL5 deficiency based on genetic testing
2. Trial and failure of at least two previous antiepileptic therapies

AGE RESTRICTION

2 years of age or older

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

ZURZUVAE

MEDICATION(S)

ZURZUVAE

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx of moderate to severe PPD
2. Major Depressive episode began in third trimester of pregnancy or within 4 weeks following delivery
3. Treatment will be initiated less than or equal to 12 months after delivery
4. Maximum duration 14 days per pregnancy

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

1 month

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

ZYDELIG

MEDICATION(S)

ZYDELIG

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. CLL dx:

a. Used in combination with rituximab

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan Year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A

ZYKADIA

MEDICATION(S)

ZYKADIA

PA INDICATION INDICATOR

1 - All FDA-Approved Indications

OFF LABEL USES

N/A

EXCLUSION CRITERIA

N/A

REQUIRED MEDICAL INFORMATION

1. Dx with metastatic non-small cell lung cancer (NSCLC) whose tumors are anaplastic lymphoma kinase (ALK)-positive
2. Patient has failed/intolerance/contraindication to Alecensa

AGE RESTRICTION

N/A

PRESCRIBER RESTRICTION

N/A

COVERAGE DURATION

Plan year

OTHER CRITERIA

N/A

PART B PREREQUISITE

N/A