

# ACNE

---

## **MEDICATION(S) SUBJECT TO STEP THERAPY**

ADAPALENE-BENZOYL PEROXIDE 0.1-2.5 % GEL, AZELEX, TRETINOIN 0.05 % GEL, TRETINOIN MICROSPHERE 0.04 % GEL, TRETINOIN MICROSPHERE 0.1 % GEL, TRETINOIN MICROSPHERE PUMP 0.04 % GEL, TRETINOIN MICROSPHERE PUMP 0.1 % GEL

## **CRITERIA**

Previous trial on at least ONE: Generic topical acne treatment

## **ADAPALENE**

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

ADAPALENE 0.1 % CREAM, ADAPALENE 0.3 % GEL, TAZAROTENE 0.1 % CREAM

### **CRITERIA**

Previous use of non-micronized tretinoin

## AJOVY

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

AJOVY

### **CRITERIA**

Previous trial on generics in two of the three following classes: a. Antidepressant: amitriptyline, nortriptyline, venlafaxine, b. Beta Blocker: atenolol, bisoprolol, metoprolol, nadolol, nebivolol, propranolol, timolol c. Anti-Epileptic agents: topiramate, valproic acid

## **ANTICONVULSANT**

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

APTIOM, BRIVIACT 10 MG TAB, BRIVIACT 10 MG/ML SOLUTION, BRIVIACT 100 MG TAB, BRIVIACT 25 MG TAB, BRIVIACT 50 MG TAB, BRIVIACT 75 MG TAB, FYCOMPA, OXCARBAZEPINE ER, SPRITAM 1000 MG TAB, SPRITAM 500 MG TAB, SPRITAM 750 MG TAB

### **CRITERIA**

Previous trial on at least TWO of the following: carbamazepine, divalproex, epilol, ethosuximide, felbamate, lamotrigine, levetiracetam tablets or solution, oxcarbazepine IR, phenytoin, tiagabine, topiramate, valproic acid, zonisamide

## **ANTIDEPRESSION**

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

APLENZIN, DRIZALMA SPRINKLE, EMSAM, FETZIMA, FETZIMA TITRATION, TRINTELLIX, VENLAFAXINE BESYLATE ER

### **CRITERIA**

Previous trial on at least TWO of the following: bupropion, citalopram, desvenlafaxine, duloxetine, escitalopram, fluoxetine, fluvoxamine, mirtazapine, paroxetine, sertraline, venlafaxine (other than 112.5mg strength)

## **ANTIPSYCHOTIC**

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

ASENAPINE MALEATE, AUVELITY, SECUADO

### **CRITERIA**

Previous trial on at least ONE of the following: aripiprazole, clozapine, fluoxetine-olanzapine, haloperidol, olanzapine, paliperidone, quetiapine, risperidone, ziprasidone

## **BELSOMRA**

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

BELSOMRA

### **CRITERIA**

Previous trial on at least ONE of the following: zolpidem IR, zolpidem ER, zaleplon, eszopiclone, temazepam, triazolam, ramelteon, trazodone

## **BEVESPI**

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

BEVESPI AEROSPHERE

### **CRITERIA**

Patient must try BOTH Anoro and Stiolto



## **BUDESONIDE**

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

BUDESONIDE ER

### **CRITERIA**

Previous trial of: Mesalamine

## CADUET

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

AMLODIPINE-ATORVASTATIN

### **CRITERIA**

Previous trial on at least ONE of the following: atorvastatin, fluvastatin, lovastatin, pravastatin, rosuvastatin, simvastatin AND Previous trial on: amlodipine

## **CLOMIPRAMINE**

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

CLOMIPRAMINE HCL, PROTRIPTYLINE HCL, TRIMIPRAMINE MALEATE

### **CRITERIA**

Previous trial on at least ONE of the following: amitriptyline, amoxapine, desipramine, doxepin, imipramine, nortriptyline

## **CORLANOR**

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

CORLANOR 5 MG/5ML SOLUTION, IVABRADINE HCL

### **CRITERIA**

Previous trial on at least ONE of the following: acebutolol, atenolol, bisoprolol, labetalol, metoprolol, nadolol, pindolol, propranolol, timolol Exception for dilated cardiomyopathy in pediatric patients

## **DAPSONE**

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

DAPSONE 5 % GEL

### **CRITERIA**

Previous use of topical clindamycin AND topical erythromycin

## DERM

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

PIMECROLIMUS

### **CRITERIA**

Previous trial on: tacrolimus AND Previous trial on at least ONE of the following: betamethasone, clobetasol, hydrocortisone, triamcinolone

## DIABETES

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

SEGLUROMET, SOLIQUA, STEGLATRO

### **CRITERIA**

Previous trial on metformin, metformin ER, metformin solution, alogliptin/metformin, glipizide/metformin, glyburide/metformin, metformin/saxagliptin, OR pioglitazone/metformin

## **DIFICID**

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

DIFICID

### **CRITERIA**

Patient has tried oral vancomycin



## **EDARBYCLOR**

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

EDARBYCLOR

### **CRITERIA**

Previous trial on at least ONE of the following: candesartan, irbesartan, losartan, olmesartan, telmisartan, valsartan, amlodipine/olmesartan, amlodipine/valsartan, candesartan/HCTZ, irbesartan/HCTZ, losartan/HCTZ, olmesartan/amlodipine/HCTZ, olmesartan/HCTZ, telmisartan/amlodipine, telmisartan/HCTZ, valsartan/HCTZ

# ENVARUSUS

---

## **MEDICATION(S) SUBJECT TO STEP THERAPY**

ASTAGRAF XL, ENVARUSUS XR

## **CRITERIA**

Patient has tried generic tacrolimus

## **EQUETRO**

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

EQUETRO

### **CRITERIA**

Previous trial on at least ONE of the following: carbamazapine ER or carbamazepine IR

## **FEMRING**

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

FEMRING

### **CRITERIA**

Previous trial on estradiol cream

## **GEMTESA**

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

GEMTESA

### **CRITERIA**

Previous trial on Myrbetriq AND one of the following: solifenacin, darifenacin, fesoterodine

## **INCRUSE**

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

INCRUSE ELLIPTA

### **CRITERIA**

Patient must try Spiriva

## **INS**

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

OMNARIS, QNASL, QNASL CHILDRENS

### **CRITERIA**

Previous trial on: fluticasone, mometasone, or flunisolide

## **MOVIPREP**

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

PLENVU

### **CRITERIA**

Previous trial on at TWO of the following: Suprep, Clenpiq, Sodium/Potas Sol Magnesium (Suprep)



## NEUAC

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

CLINDAMYCIN PHOS-BENZOYL PEROX 1-5 % GEL, CLINDAMYCIN PHOS-BENZOYL PEROX  
1.2-2.5 % GEL

### **CRITERIA**

Previous use of clindamycin-BPO 1.2%-5%

## **OLOPATADINE**

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

AZELASTINE-FLUTICASONE, OLOPATADINE HCL 0.6 % SOLUTION

### **CRITERIA**

Previous trial on: azelastine nasal spray

## **PARKINSONS**

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

RYTARY

### **CRITERIA**

Patient must try generic carbidopa/levodopa

## **PRAMIPEXOLE**

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

PRAMIPEXOLE DIHYDROCHLORIDE ER

### **CRITERIA**

Previous trial on pramipexole IR

## **QELBREE**

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

QELBREE

### **CRITERIA**

Patient has tried: atomoxetine AND one of the following: guanfacine or clonidine

## **RASUVO**

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

RASUVO

### **CRITERIA**

Previous trial on: generic methotrexate

## SOOLANTRA

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

IVERMECTIN 1 % CREAM

### **CRITERIA**

Previous trial on at least ONE of the following: Azelaic Acid or Metronidazole (topical)

# STATIN

---

## **MEDICATION(S) SUBJECT TO STEP THERAPY**

PITAVASTATIN CALCIUM

## **CRITERIA**

Previous trial on at least ONE of the following: atorvastatin, fluvastatin IR, lovastatin, pravastatin, rosuvastatin, simvastatin



## STIMULANTS

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

LISDEXAMFETAMINE DIMESYLATE 10 MG CAP, LISDEXAMFETAMINE DIMESYLATE 20 MG CAP, LISDEXAMFETAMINE DIMESYLATE 30 MG CAP, LISDEXAMFETAMINE DIMESYLATE 40 MG CAP, LISDEXAMFETAMINE DIMESYLATE 50 MG CAP, LISDEXAMFETAMINE DIMESYLATE 60 MG CAP, LISDEXAMFETAMINE DIMESYLATE 70 MG CAP, METHYLPHENIDATE

### **CRITERIA**

For lisdexamfetamine and methylphenidate patches: Previous trial on ONE of: amphetamine/dextroamphetamine, dexmethylphenidate, dextroamphetamine, or methylphenidate caps/tabs. Please note, step not required for dx of binge-eating disorder.

## **SUNOSI**

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

SUNOSI

### **CRITERIA**

Previous trial on at least one generic (e.g. modafinil, armodafinil, etc)

## **SYMLIN**

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

SYMLINPEN 120, SYMLINPEN 60

### **CRITERIA**

Previous trial on: insulin

## TAZORAC

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

TAZAROTENE 0.05 % CREAM

### **CRITERIA**

Previous use of tazarotene 0.1% cream

## TEKTURNA

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

ALISKIREN FUMARATE

### **CRITERIA**

Previous trial on at least ONE of the following: benazepril, benazepril-HCTZ, captopril, enalapril, enalapril-HCTZ, fosinopril, fosinopril-HCTZ, lisinopril, lisinopril-HCTZ, quinapril, ramipril AND

Previous trial on at least ONE of the following: candesartan, candesartan-HCTZ, irbesartan, irbesartan-HCTZ, losartan, losartan-HCTZ, olmesartan, olmesartan-HCTZ, telmisartan, telmisartan-HCTZ, valsartan, valsartan-HCTZ

## TIZANIDINE

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

TIZANIDINE HCL 2 MG CAP, TIZANIDINE HCL 4 MG CAP, TIZANIDINE HCL 6 MG CAP

### **CRITERIA**

Previous trial on: tizanidine tablets

## TRIPTAN

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

DICLOFENAC POTASSIUM(MIGRAINE), FROVATRIPTAN SUCCINATE, SUMATRIPTAN,  
ZOLMITRIPTAN 2.5 MG SOLUTION, ZOLMITRIPTAN 5 MG SOLUTION

### **CRITERIA**

Previous trial on at least ONE of the following: sumatriptan tablets/injection OR rizatriptan

## **VTAMA**

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

VTAMA

### **CRITERIA**

Previous trial on topical steroid, tretinoin, or tazarotene



## VYZULTA

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

RHOPRESSA, ROCKLATAN, VYZULTA

### **CRITERIA**

Previous trial on: latanoprost, bimatoprost, or Lumigan

## WOMENS HEALTH

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

IMVEXXY MAINTENANCE PACK, IMVEXXY STARTER PACK, PREMARIN 0.625 MG/GM CREAM

### **CRITERIA**

Previous trial on generic vaginal estradiol

## ZORYVE

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

ZORYVE

### **CRITERIA**

Previous use of BOTH the following: 1. Topical steroid, tretinoin, tazarotene 2. Vtama (Vtama not required for those under 18 years of age or those using for seborrheic dermatitis or atopic dermatitis)