



SelectHealth Medical Policies

General Medicine Policies

Table of Contents

Policy Title	Policy Number	Last Revised
Complimentary and Alternative Medicine	589	09/30/22
Long-Term Acute Care	583	07/20/16
Medical Transportation	344	09/07/22
Wound Care and Physical Therapy	469	06/20/19



COMPLEMENTARY AND ALTERNATIVE MEDICINE (CAM)

Policy # 589

Implementation Date: 1/13/17

Review Dates: 2/1/17, 12/20/18, 12/20/19, 12/14/20, 10/26/21

Revision Dates: 9/30/22

Related Medical Policies:

[#148 Medical Necessity](#)

[#241 Transcranial Magnetic Stimulation for Depression and Other Psychiatric Disorders](#)

[#252 Hippotherapy \(Equine Movement Therapy or Equine-Facilitated Psychotherapy\)](#)

[#296 Chelation Therapy](#)

[#402 Mickel Therapy for the Treatment of Fibromyalgia](#)

[#591 Computer Spinal Analysis in Chiropractic Care](#)

Disclaimer:

1. Policies are subject to change without notice.
2. Policies outline coverage determinations for SelectHealth Commercial, SelectHealth Advantage (Medicare/CMS), and SelectHealth Community Care (Medicaid/CHIP) plans. Refer to the "Policy" section for more information.

Description

Complementary and alternative medicine (CAM), also called unconventional, non-conventional, or non-traditional healthcare, is a group of diverse medical and healthcare systems, practices and products that are not typically considered to be part of traditional Western medicine. CAM assessments and therapies are proposed to reduce disease-based clinical symptoms and improve health and wellness.

Complementary medicine may be used in conjunction with Western medicine, as opposed to alternative medicine which may be used in place of Western medicine. Integrative medicine, as defined by the National Center for Complementary and Alternative Medicine (NCCAM), combines conventional medical therapies and CAM therapies for which there is scientific evidence of safety and effectiveness (NCCAM, 2015). Classifications of CAM practices include the following:

1. **Whole Medical Systems:** Whole medical systems are built upon complete systems of theory and practice. Often, these systems have evolved apart from, and earlier than, the conventional medical approach used in the United States.
2. **Biologically-Based Practices:** Biologically-based practices in CAM use substances found in nature including herbs, foods, and vitamins. Examples of these substances include dietary supplements, herbal products, and other natural products that have not been scientifically proven (e.g., using shark cartilage to treat cancer).
3. **Energy Medicine:** Energy medicine involves the use of energy fields and consist of two types of therapies:
 - a. Biofield therapies are intended to affect energy fields that purportedly surround and penetrate the human body. The existence of such fields has not yet been scientifically proven. Some forms of energy therapy are proposed to manipulate biofields by applying pressure, heat, or body manipulation.
 - b. Bioelectromagnetic-based therapies involve the unconventional use of electromagnetic fields, such as pulsed fields, magnetic fields, or alternating current or direct current fields.
4. **Manipulative and Body-Based Methods:** Manipulative and body-based methods are based on manipulation and/or movement of one or more parts of the body.
5. **Mind-Body Medicine:** Mind-body medicine uses a variety of techniques designed to enhance the mind's capacity to affect bodily function and symptoms.

General Medicine Policies, Continued

Complimentary and Alternative Medicine (CAM), continued

CAM therapies are supported by some degree of scientific evidence, but for most of the other CAM therapies key questions regarding the safety and efficacy of these therapies for specific conditions are yet to be answered through well-designed scientific studies (NCCAM, 2015).

Commercial Plan Policy/CHIP (Children's Health Insurance Program)

SelectHealth **does NOT cover complimentary or alternative medicine diagnostic testing methods, systems, therapies, or treatments because they are considered experimental, investigational, or unproven.**

SelectHealth **does NOT cover routine diagnostic testing when performed to support complementary and alternative therapies as they meet the plan's definition of investigational.**

The following is a *non-inclusive list* of excluded procedures or therapeutic interventions. This list is not intended to represent all current available testing or therapies.

acupressure and acupuncture	craniosacral therapy	laetrile
Alexander's technique	crystal healing	live blood cell analysis
Amma therapy	cupping	macrobiotics
antineoplastons	dietary supplements	meditation
antioxidant function testing (Spectrox)	ear candling	meridian therapy
aroma therapy	electromagnetic fields	Mickel therapy for fibromyalgia
art and dance therapy	equestrian therapy (hippotherapy)	mirror box therapy
auto urine therapy	essential oil therapy	moxibustion therapy
ayurvedic medicine	faith healing	MTH-68
(BELD) Bio Photonic Lymphatic Drainage Treatment	Feldenkrais therapy	music therapy
bioenergetics' analysis	Greek cancer cure test	naprapathy
biofield therapeutics	Hellerwork	Neural therapy
Bioscan	hepatic detoxification	nutrient panel testing
cellular therapy	herbal products	ozone therapy
Chelation therapy (except as outlined in med pol #296)	Homeopathic and naturopathic medicine	perineural injection therapy
chemical hair analysis	humor therapy	Pilates
Chung Moo Doe therapy	hydrogen peroxide (intravenous)	polarity therapy
martial art	hypnosis	psychodrama therapy
Coley's toxin	illimeter wave therapy	QiGong
colonic irrigation, lavage, and cleansing	immunoaugmentive therapy	Ream's testing
color therapy	inversion therapy	reflexology (zone therapy)
	iridology	Reiki
		Revici's Guided Chemo-therapy
		Rolfing
		Trager
	Kelly-Gonzales therapy	salivary hormone panels

General Medicine Policies, Continued

Complimentary and Alternative Medicine, continued

Telomere testing	Traditional Chinese	Vitamin therapy in the
therapeutic touch	medicine	absence of a documented
Tichuris suis ova therapy	Tui Na	deficiency
		yoga

SelectHealth Advantage (Medicare/CMS)

Coverage is determined by the Centers for Medicare and Medicaid Services (CMS); if a coverage determination has not been adopted by CMS, and InterQual criteria are not available, the SelectHealth Commercial policy applies. For the most up-to-date Medicare policies and coverage, please visit their search website <http://www.cms.gov/medicare-coverage-database/overview-and-quick-search.aspx?from2=search1.asp&> or [the manual website](#)

SelectHealth Community Care (Medicaid)

Coverage is determined by the State of Utah Medicaid program; if Utah State Medicaid has no published coverage position and InterQual criteria are not available, the SelectHealth Commercial criteria will apply. For the most up-to-date Medicaid policies and coverage, please visit their website <http://health.utah.gov/medicaid/manuals/directory.php> or the [Utah Medicaid code Look-Up tool](#)

Summary of Medical Information

The Federal Food and Drug Act of 1906, The Wiley Act, empowers the FDA Center for Food Safety and Nutrition to remove unsafe food substances and botanicals from the market, and gives the FDA regulatory oversight for substances added to food, including monitoring safe use. The FDA maintains that a drug is any substance or mixture of substances intended for the cure, mitigation, diagnosis, or prevention of disease (FDA, 2009).

Dietary supplements are regulated differently than prescription and over-the-counter drug products. Manufacturers of dietary supplements are responsible for ensuring that their products are safe. While the FDA monitors adverse effects after dietary supplement products are on the market, newly marketed dietary supplements are not subject to premarket approval or a specific post-market surveillance period. Per the Dietary Supplement Health and Education Act of 1994 (DSHEA), the burden of proof rests on the FDA to show that a product is unsafe. Manufacturers are not required to submit substantiation of benefit data to the FDA. The Federal Trade Commission (FTC) is charged with accurate marketing and advertising claims.

According to the FDA, dietary supplements in today's market include one or a combination of: vitamins, minerals, herbals, botanicals, amino acids, any dietary substance used to supplement the diet by increasing total dietary intake, and a concentrate, metabolite, constituent, or extract. The FDA states that, while some supplements may help ensure that the individual consumes adequate amounts of essential nutrients needed for optimal health and performance, dietary supplements cannot be promoted as a treatment or a cure.

In December 2006, the FDA issued a draft guidance document for the regulation of CAM products. The draft was issued because increased use of CAM in the United States has caused confusion regarding which products are subject to regulation under the Federal Food, Drug, and Cosmetic Act (Act) or the Public Health Service Act (PHS Act) and because the number of CAM products being imported into the United States has increased. The document provides guidance as to when a CAM product is subject to the Act or the PHS Act. The FDA cites the NCCAM's definition and categories of CAM in the draft. According to the new guidance, if the labeling of a dietary supplement includes the term "to treat," that supplement will be regulated as a drug under the Act. Biological products (e.g., virus, therapeutic serum, toxin, antitoxin, vaccine) will be regulated under the PHS Act (FDA, 2010; FDA, 2007).

Several systematic reviews have evaluated multiple CAM therapies for the treatment of various conditions including asthma, cancer, depression, diabetes, hypertension, irritable bowel syndrome, pain management, psoriasis, Raynaud's phenomenon, rheumatoid arthritis, and rhinitis. Although some

General Medicine Policies, Continued

Complimentary and Alternative Medicine (CAM), continued

studies reported clinical improvement with some modalities, overall, the authors agreed that there is insufficient evidence to support CAM for the treatment of these conditions. Studies are limited by small patient populations, minimal and short-term follow-ups, variability in dosage and unknown quality of oral supplements, few evaluations of side effects, inconsistent and inconclusive outcomes, and no controls or comparisons to traditional Western medical therapies.

American Academy of Allergy, Asthma & Immunology (AAAAI): In their clinical review of CAM (Mainardi, et al., 2009) which included vitamins D, E, C and A, magnolol, quercetin, resveratrol, ma huang (ephedrine sinica), Ayurvedic medicine, Kampo medicine for the treatment of asthma, atopic dermatitis, and allergic rhinitis, the AAAAI concluded that further studies are needed using larger sample sizes, longer study durations, comparable absolute measures, and well-constructed study designs that control for biases. They also stated that the following are unknown: the true efficacy and safety of CAM therapies, the efficacy of CAM therapies alone (as alternatives) in the treatment of various disorders, the individual CAM therapeutic mechanism of effects (some may be multiple), the active component of individual CAM therapies, the potential drug-drug and drug-herb-phytochemical and vitamin interactions.

American Academy of Neurology (AAN): AAN (2014) conducted a systematic review of the literature to develop recommendations for CAM for the treatment of multiple sclerosis. Due to the lack of evidence or the poor quality of the evidence, AAN concluded that the evidence was insufficient to support or refute the use of Chinese medicine, hypnotherapy, massage therapy, hypnosis, mindfulness training, music therapy, naturopathic medicine, neural therapy, progressive muscle relaxation, tai chi, and yoga. Based on available studies, AAN concluded the following: 1) Ginkgo biloba is ineffective for improving cognitive function but possibly effective in reducing fatigue (level A); 2) low-fat diet with omega-3 fatty acid supplement is probably ineffective for reducing MS-related relapse, disability, MRI lesions, or for improving fatigue or quality of life (QOL) (level B); 3) reflexology is possibly effective for reducing MS-associated paresthesia but there is a lack of data to support or refute this modality for pain, health-related QOL, disability, spasticity, fatigue, cognition, bowel/bladder function, depression, anxiety, or insomnia (level C); 4) Bee sting therapy is possibly ineffective (level C); 5) Magnetic therapy is probably effective for reducing fatigue (level B), but not depression (level B); 6) Safety and efficacy of all other CAM remains unknown (level 4).

American Academy of Pediatrics (AAP): The 2017 AAP Task Force on Complementary and Alternative Medicine, the Provisional Section on Complementary, Holistic, and Integrative Medicine published guidance on the use of CAM in pediatrics. The Task Force concluded that pediatricians and other clinicians who care for children have the responsibility to advise and counsel patients about relevant, safe, effective, and age-appropriate health therapies including CAM and should routinely inquire as to whether or not the patient is using any specific CAM therapies. They advised the clinician to work with the parents to consider and evaluate all appropriate treatments and monitor the patient's response to treatments. They also stated that the physician should be knowledgeable about CAM therapies and evidence-based information.

American Cancer Society (ACS): In their operational statement on CAM methods for cancer management, the ACS (2018) urged patients who are thinking about using complementary or non-mainstream therapies to discuss them with their healthcare team first. ACS noted that complementary methods that may be helpful and safe include aromatherapy, art therapy, massage therapy, meditation, music therapy, prayer and spirituality, tai chi, and yoga. Although some therapies are safe and may be helpful, some CAM therapies have been associated with serious problems and death.

American College of Chest Physicians (ACCP): The ACCP (Gabay, et al., 2017) published evidence-based clinical practice guidelines on complementary therapies and integrative medicine. Despite no evidence for efficacy, there is widespread use. The use of mind-body modalities as part of a multidisciplinary approach to treating the symptoms of cancer-related pain, nausea, and vomiting associated with chemotherapy, anxiety, and sleep and mood disturbances may be helpful. Yoga and massage therapy may be beneficial in reducing fatigue, anxiety, and/or pain.

American College of Rheumatology (ACR): The ACR (2012) position statement on complementary and alternative medicine for rheumatic diseases supports the integration of CAM modalities "proven to be safe and effective by scientifically rigorous clinical trials published in the biomedical peer review literature" and advised caution in using those therapies not scientifically studied. For interventions for which randomized controlled trials are not feasible: "Innovative methods of evaluation are needed, as are measures and standards for the generation and interpretation of evidence."

American Psychiatric Association: The American Psychiatric Association's Task Force on Complementary and Alternative Medicine (Freeman, et al., 2010) conducted a systematic review of

General Medicine Policies, Continued

Complimentary and Alternative Medicine (CAM), continued

randomized controlled trials to evaluate the evidence on commonly used CAM therapies for the treatment of major depressive disorder (MDD). Therapies included omega-3 fatty acids, St. John's wort (Hypericum), folate, S-adenosyl-L-methionine (SAMe), bright light therapy, exercise, and mindfulness psychotherapies (i.e., mindfulness-based cognitive therapy, problem-solving therapy, well-being therapy). The Task Force concluded that although some CAM therapies were promising, more rigorous studies to determine their role in the treatment of MDD were necessary. It was noted that the greatest risk of pursuing a CAM therapy is the possible delay of other well-established treatments.

National Cancer Institute (NCI): NCI (2020) states that cancer patients using or considering complementary or alternative therapy should discuss this decision with their healthcare provider to ensure coordination of care. NCI notes that some complementary and alternative therapies may interfere with standard treatment or may be harmful when used with conventional treatment. Patients should become informed about the therapy, including whether the results of scientific studies support the claims that are made for it.

National Comprehensive Cancer Network (NCCN): In their Clinical Practice Guidelines on Cancer-Related Fatigue, NCCN (2016) stated that: "Complementary therapies including massage therapy, yoga, muscle relaxation, and stress reduction based on mindfulness have been evaluated in some studies and the data suggested that these therapies might be effective in reducing fatigue in cancer patients." NCCN (2016) listed imagery, hypnosis, distraction training, and relaxation training as nonpharmacological coping skills for the treatment of adult cancer pain. Relaxation/systemic desensitization, hypnosis/guided imagery and music therapy are noted as interventions for anticipatory nausea and vomiting (NCCN, 2016).

Society for Integrative Oncology (SIO): The SIO (Deng, et al., 2013) evidence-based practice guidelines included a systematic review of complementary therapy and botanicals in the care of cancer patients. Mind-body modalities, bioenergy field therapy, and dietary supplements were evaluated. SIO recommendations included the use of massage, yoga and mind-body modalities may be helpful in a multidisciplinary approach to reduce anxiety, mood disturbance, sleep disturbance, pain, and anticipatory chemotherapy-induced nausea and vomiting, and to improve overall quality of life. The SIO noted that in general, clinical trials of CAM therapies suffer from limitations in design and implementation of studies.

In 2014, SIO conducted a systematic review of the literature to develop guidelines on integrative therapies for supportive care for breast cancer patients. The recommendations included the use of music therapy, meditation, yoga, message and/or relaxation for the relief of anxiety, stress, depression/mood and/or fatigue. In addition, Qigong, mistletoe, and reflexology were recommended as supportive activities for quality of life and physical functioning. Grade A recommendations (high certainty that the net benefit is substantial) were given to meditation, relaxation, and yoga for the treatment of depression/mood and for quality of life and physical functioning. The remaining recommendations were lower grades. Forty of the 53 recommendations were rated a C (moderate certainty that the net benefit is small), D (moderate/high certainty that modality has no benefit or H (moderate certainty that harms outweigh benefits). Limitations of the literature included lack of standardization of intervention and the variety of settings in which the interventions were used (Greenlee, et al., 2014).

Outside the United States

According to the World Health Organization (WHO) traditional medicine (TM) is either the mainstay of healthcare delivery or serves as a complement to it. In some countries, traditional medicine or non-conventional medicine are termed complementary medicine (CM). Tradition and complementary medicine (T&CM) is found in almost every country in the world and has a long history of use in health maintenance and in disease prevention and treatment, especially in the treatment of chronic disease. The use of T&CM varies by country based on culture and accessibility. T&CM products include herbs, herbal materials, herbal preparations, and finished herbal products. T&CM practices include Ayurveda, traditional Chinese medicine, qigong, tai chi, yoga, thermal medicine, and other physical, mental, spiritual and mind-body therapies. Halotherapy, or salt therapy, has been used in Europe for over 20 years. In a few countries, certain types of T&CM have been completely integrated into the healthcare system. For example, in China, traditional Chinese medicine and conventional medicine are practiced alongside each other at every level of the healthcare service. By the latter half of the 19th century, homeopathy was practiced throughout Europe, Asia, and North America. Homeopathy has been integrated into the national healthcare systems of India, Mexico, Pakistan, Sri Lanka, and the United Kingdom. The regulation of TC&M varies from country to country (WHO, 2014; WHO 2001).

National Institute for Clinical Excellence (NICE): NICE (United Kingdom) (2015) published a guideline document on the treatment of irritable bowel syndrome (IBS) in adults. The review included the use of homeopathic medicine, Chinese herbal medicine, and reflexology. Regarding homeopathy for IBS, NICE

General Medicine Policies, Continued

Complimentary and Alternative Medicine (CAM), continued

stated that randomized trials for the past 30 years were not found. Only one quasi-randomized trial was found regarding the use of reflexology for the treatment of IBS (n=34). Six trials met inclusion criteria for evaluation of the use of Chinese herbal medicines. The studies utilized various combinations of herbal preparations. The Guideline Development Group (GDG) concluded that the review of evidence suggests that some herbal preparations may be clinically effective in people with IBS and are well-tolerated. However, the GDG believed there were too many uncertainties regarding type and dose of herbal medicines to make a recommendation for practice and proposed that these interventions should be investigated further in a research recommendation.

New Zealand Guidelines Group: In a mental health disorders guideline for the management of depression, The New Zealand Guidelines Group (2008) conducted a systematic review of randomized controlled trials of therapies used for the treatment of depression in adults in primary care. The authors stated that: "There was insufficient evidence to determine whether any complementary or alternative medicines are effective for the treatment of depression in young people." Very few randomized controlled trials were found, and none were found for the use of St. John's Wort for depression in young people. The groups also pointed out that due to safety concerns, patients using St. John's wort should be advised of possible drug interactions. One small study (n=28) supported the possibility that omega-3 supplements may be effective in the treatment of childhood depression and it is proposed that omega-3 may be useful in the treatment of women in antenatal or postnatal periods, but no controlled trials were available.

Royal Australian and New Zealand College of Obstetrician and Gynecologists (RANZCOG): In their statement on the use of vitamin and mineral supplements during pregnancy, the RANZCOG (2014) stated that there is a lack of high-quality evidence to support the use of omega-3 fatty acid supplements during pregnancy. However, their consensus-based recommendation is that women with a low dietary intake of Omega-3 fatty acids should consider using a dietary supplement.

Complementary and alternative medicine (CAM) encompasses a wide array of practices and treatment modalities that deviate from conventional Western medical treatment. They may be used in lieu of, or in conjunction with, traditional medical therapies. Overall, randomized controlled trials included heterogeneous small patient populations, short-term follow-ups, various controls, treatment regimens and outcome measures, inconsistent and conflicting outcomes, and poor methodology. Some CAM testing methods and therapies/treatments lack standardization of regimens and/or practitioner training. Systematic reviews have been unable to make firm conclusions about CAM testing methods and therapies due to the study limitations, and in some cases, lack of data. The evidence in the published peer-reviewed scientific literature does not support the safety, efficacy, and/or clinical utility of the diagnostic testing and therapies discussed.

Billing/Coding Information

CPT CODES

20550	Injection(s); single tendon sheath, or ligament, aponeurosis (eg, plantar "fascia")
45399	Unlisted procedure, colon
64415	Injection, anesthetic agent; brachial plexus, single
64418	Injection, anesthetic agent; suprascapular nerve
64450	Injection, anesthetic agent; other peripheral nerve or branch
84999	Unlisted chemistry procedure
86353	Lymphocyte transformation, mitogen (phytomitogen) or antigen induced blastogenesis
86849	Unlisted immunology procedure
90880	Hypnotherapy
90899	Unlisted psychiatric service or procedure
96372	Therapeutic, prophylactic, or diagnostic injection (specify substance or drug); subcutaneous or intramuscular
96379	Unlisted therapeutic, prophylactic, or diagnostic intravenous or intra-arterial injection or infusion
96549	Unlisted chemotherapy procedure
97139	Unlisted therapeutic procedure (specify)

General Medicine Policies, Continued

Complimentary and Alternative Medicine (CAM), continued

97799	Unlisted physical medicine/rehabilitation service or procedure
99199	Unlisted special service, procedure or report

HCPSC CODES

P2031	Hair analysis (excluding arsenic)
S3650	Saliva test, hormone level; during menopause
S3652	Saliva test, hormone level; to assess preterm labor risk
M0075	Cellular therapy
M0076	Prolotherapy
J3570	Laetrile, amygdalin, vitamin B-17
A9152	Single vitamin/mineral/trace element, oral, per dose, not otherwise specified
A9153	Multiple vitamins, with or without minerals and trace elements, oral, per dose, not otherwise specified
S9451	Exercise classes, nonphysician provider, per session
G0176	Activity therapy, such as music, dance, art or play therapies not for recreation, related to the care and treatment of patient's disabling mental health problems, per session (45 minutes or more)
H2032	Activity therapy, per 15 minutes

Key References

1. AACT, A. (2015). "List of Commonly Used Tests and Treatments to Question." Accessed December 14, 2020. Retrieved Sep 5, 2017, from <http://www.choosingwisely.org/toxicology-societies-release-list-of-commonly-used-tests-and-treatments-to-question-as-part-of-choosing-wisely-campaign-2/>.
2. Al-Harasi S, Ashley PF, Moles DR, Parekh S, Walters V. Hypnosis for children undergoing dental treatment. Cochrane Database of Systematic Reviews 2010, Issue 8. Art. No.: CD007154. DOI: 10.1002/14651858.CD007154.pub2.
3. Alraek T, Lee MS, Choi TY, Cao H, Liu J. Complementary and alternative medicine for patients with chronic fatigue syndrome: a systematic review. BMC Complement Altern Med. 2011 Oct 7; 11:87.
4. American Academy of Neurology (AAN): Summary of evidence-based guideline: complementary and alternative medicine in multiple sclerosis. Mar 2014. Accessed Jun 5, 2015. Available at URL address: <https://www.aan.com/Guidelines/home/ByTopic?topicId=18>
5. American Cancer Society. Complementary and alternative medicine. Accessed December 14, 2020. Available at URL address: <http://www.cancer.org/Treatment/TreatmentsandSideEffects/ComplementaryandAlternativeMedicine/complementary-and-alternative-methods-for-cancer-management>
6. American College of Rheumatology. Position statement. Complementary and alternative medicine for rheumatic diseases. Aug 2008. Accessed December 14, 2020. Available at URL address: <http://www.rheumatology.org/practice/clinical/position/index.asp>
7. American Naprapathic Association. 2020. Accessed December 14, 2020. Available at URL address: <http://www.naprapathy.org/>
8. Babbar S, Parks-Savage AC, Chauhan SP. Yoga during pregnancy: a review. Am J Perinatol. 2012 Jun;29(6):459-64.
9. Bager P, Amved J, Rønberg S, Wohlfahrt J, Poulsen LK, Westergaard T, Petersen HW, Kristensen B, Thamsborg S, Roepstorff A, Kapel C, Melbye M. Trichuris suis ova therapy for allergic rhinitis: a randomized, double-blind, placebo-controlled clinical trial. J Allergy Clin Immunol. 2010 Jan;125(1):123-30 e1-3.
10. Balíková M. Hair analysis for drugs of abuse. Plausibility of interpretation. Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub. 2005 Dec;149(2):199-207.
11. Bardia A, Barton DL, Prokop LJ, Bauer BA, Moynihan TJ. Efficacy of complementary and alternative medicine therapies in relieving cancer pain: a systematic review. J Clin Oncol. 2006 Dec 1;24(34):5457-64.
12. Bennett C, Underdown A, Barlow J. Massage for promoting mental and physical health in typically developing infants under the age of six months. Cochrane Database of Systematic Reviews 2013, Issue 4. Art. No.: CD005038. DOI: 10.1002/14651858.CD005038.pub3.
13. Bernstein IL, Li JT, Bernstein DI, Hamilton R, Spector SL, Tan R, Sicherer S, Golden DB, Khan DA, Nicklas RA, Portnoy JM, Blessing-Moore J, Cox L, Lang DM, Oppenheimer J, Randolph CC, Schuller DE, Tilles SA, Wallace DV, Levitin E, Weber R, American Academy of Allergy, Asthma and Immunology, American College of Allergy, Asthma and Immunology. Allergy diagnostic testing: an updated practice parameter. Part 1. Ann Allergy Asthma Immunol 2008 Mar;100(3 Suppl 3): S15-S66. Accessed Jun 4, 2015. Available at URL address: <http://www.aaaai.org/practice-resources/statements-and-practice-parameters/practice-parameter-guidelines.aspx>
14. Bronson C, Brewerton K, Ong J, Palanca C, Sullivan SJ. Does hippotherapy improve balance in persons with multiple sclerosis: a systematic review. Eur J Phys Rehabil Med. 2010 Sep;46(3):347-53.
15. Büssing A, Ostermann T, Lüdtker R, Michalsen A. Effects of yoga interventions on pain and pain-associated disability: a meta-analysis. J Pain. 2012 Jan;13(1):1-9.
16. Cacchio A, De Blasis E, De Blasis V, Santilli V, Spacca G. Mirror therapy in complex regional pain syndrome type 1 of the upper limb in stroke patients. Neurorehabil Neural Repair. 2009 Oct;23(8):792-9.
17. Candy B, Jones L, Varagunam M, Speck P, Tookman A, King M. Spiritual and religious interventions for well-being of adults in the terminal phase of disease. Cochrane Database of Systematic Reviews 2012, Issue 5. Art. No.: CD007544. DOI:

General Medicine Policies, Continued

Complimentary and Alternative Medicine (CAM), continued

18. Cao H, Han M, Li X, Dong S, Shang Y, Wang Q, Xu S, Liu J. Clinical research evidence of cupping therapy in China: a systematic literature review. *BMC Complement Altern Med*. 2010 Nov 16; 10:70.
19. Cao H, Li X, Liu J. An updated review of the efficacy of cupping therapy. *PLoS One*. 2012;7(2): e31793. doi: 10.1371/journal.pone.0031793.
20. Caprara DL, Klein J, Koren G. Diagnosis of fetal alcohol spectrum disorder (FASD): fatty acid ethyl esters and neonatal hair analysis. *Ann Ist Super Sanita*. 2006;42(1):39-45.
21. Carlson JJ, Farquhar JW, DiNucci E, Ausserer L, Zehnder J, Miller D, Berra, Hagerty L, Haskell WL. Safety and efficacy of a ginkgo biloba-containing dietary supplement on cognitive function, quality of life, and platelet function in healthy, cognitively intact older adults. *J Am Diet Assoc*. 2007 Mar;107(3):422-32.
22. Cassileth BR, Deng GE, Gomez JE, Johnstone PA, Kumar N, Vickers AJ; American College of Chest Physicians. Complementary therapies and integrative oncology in lung cancer: ACCP evidence-based clinical practice guidelines (2nd edition). *Chest*. 2007 Sep;132(3 Suppl):340S-354S.
23. Cepeda MS, Carr DB, Lau J, Alvarez H. Music for pain relief. *Cochrane Database of Systematic Reviews* 2006, Issue 2. Updated 2010. Art. No.: CD004843. DOI: 10.1002/14651858.CD004843.pub2.
24. Chan CL, Wang CW, Ho RT, Ng SM, Chan JS, Ziea ET, Wong VC. A systematic review of the effectiveness of qigong exercise in supportive cancer care. *Support Care Cancer*. 2012 Jun;20(6):1121-33. doi: 10.1007/s00520-011-1378-3.
25. Choi TY, Choi J, Kim KH, Lee MS. Moxibustion for the treatment of osteoarthritis: a systematic review and meta-analysis. *Rheumatol Int*. 2012 Oct;32(10):2969-78.
26. Clement K, Covertson CR, Johnson MJ, Dearing K. St. John's wort and the treatment of mild to moderate depression: a systematic review. *Holist Nurs Pract*. 2006 Jul-Aug;20(4):197-203.
27. Cooley K, Szczurko O, Perri D, Mills EJ, Bernhardt B, Zhou Q, Seely D. Naturopathic care for anxiety: a randomized controlled trial ISRCTN78958974. *PLoS One*. 2009 Aug 31;4(8):e6628.
28. Crawford MJ, Killaspy H, Barnes TR, Barrett B, Byford S, Clayton K, Dinsmore J, Floyd S, Hooley A, Johnson T, Kalitzi E, King M, Leurent B, Maratos A, O'Neill FA, Osborn DP, Patterson S, Soteriou T, Tyrer P, Waller D; MATISSE project team. Group art therapy as an adjunctive treatment for people with schizophrenia: multicentre pragmatic randomised trial. *BMJ*. 2012 Feb 28; 344: e846. doi: 10.1136/bmj.e846.
29. Davidson JR, Crawford C, Ives JA, Jonas WB. Homeopathic treatments in psychiatry: a systematic review of randomized placebo-controlled studies. *J Clin Psychiatry*. 2011 Jun;72(6):795-805.
30. Deng GE, Rausch SM, Jones LW, Gulati A, Kumar NB, Greenlee H, Pietanza MC, Cassileth BR. Complementary therapies and integrative medicine in lung cancer: Diagnosis and management of lung cancer, 3rd ed: American College of Chest Physicians evidence-based clinical practice guidelines. *Chest*. 2013 May;143(5 Suppl):e420S-36S. doi: 10.1378/chest.12-2364. Accessed Jun 4, 2015. Available at URL address: <https://www.integrativeonc.org/integrative-oncology-guidelines>
31. Ding M. Tai Chi for stroke rehabilitation: a focused review. *Am J Phys Med Rehabil*. 2012 Dec;91(12):1091-6.
32. Ebrahim Soltani A, Mohammadinasab H, Goudarzi M, Arbabi S, Mohtaram R, Afkham K, Momenzadeh S, Darabi ME. Acupressure using ondansetron versus metoclopramide on reduction of postoperative nausea and vomiting after strabismus surgery. *Arch Iran Med*. 2010 Jul;13(4):288-93.
33. Ernst E, Pittler MH, Wider B, Boddy K. Complementary/alternative medicine for supportive cancer care: development of the evidence-base. *Support Care Cancer*. 2007 May;15(5):565-8.
34. Ernst E, Posadzki P, Lee MS. Reflexology: an update of a systematic review of randomised clinical trials. *Maturitas*. 2011 Feb;68(2):116-20.
35. Ernst E, Schmidt K, Baum M. Complementary/Alternative therapies for the treatment of breast cancer. A systematic review of randomized clinical trials and a critique of current terminology. *Breast J*. 2006 Nov-Dec;12(6):526-30.
36. Ezendam D, Bongers RM, Jannink MJ. Systematic review of the effectiveness of mirror therapy in upper extremity function. *Disabil Rehabil*. 2009;31(26):2135-49.
37. Filipek PA, Accardo PJ, Ashwal S, Baranek GT, Cook EH Jr, Dawson G, Gordon B, Gravel JS, Johnson CP, Kallen RJ, Levy SE, Minshew NJ, Ozonoff S, Prizant BM, Rapin I, Rogers SJ, Stone WL, Teplin SW, Tuchman RF, Volkmar FR. Practice parameter: screening and diagnosis of autism: report of the Quality Standards Subcommittee of the American Academy of Neurology and the Child Neurology Society. *Neurology*. 2000 Aug 22;55(4):468-79. Posted Mar 26, 2013. Accessed December 14, 2020. Available at URL address: <http://www.childneurologysociety.org/resources/resources-detail-view/practice-parameter-screening-and-diagnosis-of-autism>
38. Flower A, Liu JP, Lewith G, Little P, Li Q. Chinese herbal medicine for endometriosis. *Cochrane Database of Systematic Reviews* 2012, Issue 5. Art. No.: CD006568. DOI: 10.1002/14651858.CD006568.pub3
39. Flyckt RL, Liu J, Frasure H, Wekselman K, Buch A, Kingsberg SA. Comparison of salivary versus serum testosterone levels in postmenopausal women receiving transdermal testosterone supplementation versus placebo. *Menopause*. 2009 Jul-Aug;16(4):680-8.
40. Freeman MP, Fava M, Lake J, Trivedi MH, Wisner KL, Mischoulon D. Complementary and alternative medicine in major depressive disorder: the American Psychiatric Association Task Force report. *J Clin Psychiatry*. 2010 Jun;71(6):669-81.
41. Gagnier JJ, DeMelo J, Boon H, Rochon P, Bombardier C. Quality of reporting of randomized controlled trials of herbal medicine interventions. *Am J Med*. 2006 Sep;119(9):800e1-11.
42. Genova Diagnostics. ION (Nutritional Test) Accessed December 14, 2020. Available at URL address: <http://www.gdx.net/product/10157>
43. Greenlee H1, Balneaves LG2, Carlson LE2, Cohen M2, Deng G2, Hershman D2, Mumber M2, Perlmutter J2, Seely D2, Sen A2, Zick SM2, Tripathy D2; Society for Integrative Oncology Guidelines Working Group. Clinical practice guidelines on the use of integrative therapies as supportive care in patients treated for breast cancer. *J Natl Cancer Inst Monogr*. 2014 Nov;2014(50):346-58. Accessed December 14, 2020. Available at URL address: <https://www.integrativeonc.org/integrative-oncology-guidelines>
44. Guo Z, Jia X, Liu JP, Liao J, Yang Y. Herbal medicines for advanced colorectal cancer. *Cochrane Database of Systematic Reviews* 2012, Issue 5. Art. No.: CD004653. DOI: 10.1002/14651858.CD004653.pub2.
45. Harder H, Parlour L, Jenkins V. Randomised controlled trials of yoga interventions for women with breast cancer: a systematic literature review. *Support Care Cancer*. 2012 Dec;20(12):3055-64.
46. Hilton MP, Zimmermann EF, Hunt WT. Ginkgo biloba for tinnitus. *Cochrane Database of Systematic Reviews* 2013, Issue 3. Art. No.: CD003852. DOI:
47. IntraCellular Diagnostics, Inc. ©. ExaTest. 2014. Accessed Jun 4, 2015. Available at URL address: <http://www.exatest.com/>
48. Jain S, Mills PJ. Biofield therapies: helpful or full of hype? A best evidence synthesis. *Int J Behav Med*. 2010 Mar;17(1):1-16.

General Medicine Policies, Continued

Complimentary and Alternative Medicine (CAM), continued

49. Jing Z, Yang X, Ismail KMK, Chen XY, Wu T. Chinese herbal medicine for premenstrual syndrome. *Cochrane Database of Systematic Reviews* 2009, Issue 1. Art. No.: CD006414. DOI: 10.1002/14651858.CD006414.pub2.
50. Jorm AF, Allen NB, O'Donnell CP, Parslow RA, Purcell R, Morgan AJ. Effectiveness of complementary and self-help treatments for depression in children and adolescents. *Med J Aust.* 2006 Oct 2;185(7):368-72.
51. Karsch-Völk M, Barrett B, Kiefer D, Bauer R, Ardjomand-Woelkart K, Linde K. Echinacea for preventing and treating the common cold. *Cochrane Database of Systematic Reviews* 2014, Issue 2. Art. No.: CD000530. DOI: 10.1002/14651858.CD000530.pub3.
52. Kemper KJ, Vohra S, Walls R; Task Force on Complementary and Alternative Medicine; Provisional Section on Complementary, Holistic, and Integrative Medicine. American Academy of Pediatrics. The use of complementary and alternative medicine in pediatrics. *Pediatrics.* 2008 Dec;122(6):1374-86.
53. Kim JI, Choi JY, Lee H, Lee MS, Ernst E. Moxibustion for hypertension: a systematic review. *BMC Cardiovasc Disord.* 2010 Jul 5; 10:33.
54. Lakhani SE, Vieira KF. Nutritional and herbal supplements for anxiety and anxiety-related disorders: systematic review. *Nutr J.* 2010 Oct 7; 9:42.
55. Lau HLC, Kwong JSW, Yeung F, Chau PH, Woo J. Yoga for secondary prevention of coronary heart disease. *Cochrane Database of Systematic Reviews* 2012, Issue 12. Art. No.: CD009506. DOI: 10.1002/14651858.CD009506.pub2.
56. Lee MS, Choi TY, Park JE, Lee SS, Ernst E. Moxibustion for cancer care: a systematic review and meta-analysis. *BMC Cancer.* 2010c Apr 7; 10:130.
57. Lee DH, Kim JI, Lee MS, Choi TY, Choi SM, Ernst E. Moxibustion for ulcerative colitis: a systematic review and meta-analysis. *BMC Gastroenterol.* 2010a Apr 7; 10:36.
58. Lee MS, Shin BC, Kim JI, Han CH, Ernst E. Moxibustion for stroke rehabilitation: systematic review. *Stroke.* 2010b Apr;41(4):817-20.
59. Li AW, Goldsmith CA. The effects of yoga on anxiety and stress. *Altern Med Rev.* 2012 Mar;17(1):21-35.
60. Li L, Dou L, Leung PC, Wang CC. Chinese herbal medicines for threatened miscarriage. *Cochrane Database of Systematic Reviews* 2012, Issue 5. Art. No.: CD008510. DOI: 10.1002/14651858.CD008510.pub2.
61. Lim EC, Poh RL, Low AY, Wong WP. Effects of Pilates-based exercises on pain and disability in individuals with persistent nonspecific low back pain: a systematic review with meta-analysis. *J Orthop Sports Phys Ther.* 2011 Feb;41(2):70-80.
62. Madden K, Middleton P, Cyna AM, Matthewson M, Jones L. Hypnosis for pain management during labour and childbirth. *Cochrane Database of Systematic Reviews* 2012, Issue 11. Art. No.: CD009356. DOI: 10.1002/14651858.CD009356.pub2.
63. Magalhaes FN, Dotta L, Sasse A, Teixeira MJ, Fonoff ET. Ozone therapy as a treatment for low back pain secondary to herniated disc: a systematic review and meta-analysis of randomized controlled trials. *Pain Physician.* 2012 Mar-Apr;15(2): E115-29.
64. Mainardi T, Kapoor S, Bielory L. Complementary and alternative medicine: herbs, phytochemicals and vitamins and their immunologic effects. *J Allergy Clin Immunol.* 2009 Feb;123(2):283-94.
65. Matthews A, Haas DM, O'Mathúna DP, Dowswell T, Doyle M. Interventions for nausea and vomiting in early pregnancy. *Cochrane Database of Systematic Reviews* 2014, Issue 3. Art. No.: CD007575. DOI: 10.1002/14651858.CD007575.pub3.
66. Meeks TW, Wetherell JL, Irwin MR, Redwine LS, Jeste DV. Complementary and alternative treatments for late-life depression, anxiety, and sleep disturbance: a review of randomized controlled trials. *J Clin Psychiatry.* 2007 Oct;68(10):1461-71.
67. Miyasaki JM, Shannon K, Voon V, Ravina B, Kleiner-Fisman G, Anderson K, Shulman LM, Gronseth G, Weiner WJ; Quality Standards Subcommittee of the American Academy of Neurology. Practice Parameter: evaluation and treatment of depression, psychosis, and dementia in Parkinson disease (an evidence-based review): report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology.* 2006 Apr 11;66(7):996-1002.
68. Morgan AJ, Jorm AF. Self-help interventions for depressive disorders and depressive symptoms: a systematic review. *Ann Gen Psychiatry.* 2008 Aug 19; 7:13.
69. Moseley GL, Gallace A, Spence C. Is mirror therapy all it is cracked up to be? Current evidence and future directions. *Pain.* 2008 Aug 15;138(1):7-10.
70. Moser HW, Raymond GV, Lu SE, Muenz LR, Moser AB, Xu J, Jones RO, Loes DJ, Melhem ER, Dubey P, Bezman L, Breerton NH, Odone A. Follow-up of 89 asymptomatic patients with adrenoleukodystrophy treated with Lorenzo's oil. *Arch Neurol.* 2005 Jul;62(7):1073-80.
71. Myers SM, Johnson CP; American Academy of Pediatrics Council on Children With Disabilities. Management of children with autism spectrum disorders. *Pediatrics.* 2007 Nov;120(5):1162-82.
72. Nahas R, Moher M. Complementary and alternative medicine for the treatment of type 2 diabetes. *Can Fam Physician.* 2009 Jun;55(6):591-6.
73. National Cancer Institute (NCI). Complementary and alternative medicine in cancer treatment Nov 4, 2020. Accessed December 14, 2020. Available at URL address: <http://www.cancer.gov/about-cancer/treatment/cam>
74. National Center for Complementary and Alternative Medicine. Health Topics A-Z. Accessed December 14, 2020. Available at URL address: <http://nccam.nih.gov/health/atoz.htm>
75. National Center for Complementary and Alternative Medicine (NCCAM). Complementary, Alternative, or Integrative Health: What's In a Name? July 2018. Accessed December 14, 2020. Available at URL address: <http://nccam.nih.gov/health/whatiscam/>
76. National Comprehensive Cancer Network® (NCCN®). NCCN clinical practice guidelines in oncology™. Adult cancer pain. V.2.2017. May 18, 2016 Accessed December 14, 2020. Available at URL address: <https://www.nccn.org/about/news/ebulletin/ebulletindetail.aspx?bulletinid=1582>
77. National Comprehensive Cancer Network® (NCCN®). NCCN clinical practice guidelines in oncology™. Antiemesis. V.1.2016. Apr 1, 2015. Accessed December 14, 2020. Available at URL address: <https://www.nccn.org/patients/guidelines/content/PDF/nausea-patient.pdf82>
78. National Institute for Health and Clinical Excellence (NICE). National clinical practice guideline CG61. Irritable bowel syndrome: full guidance. Feb 2015. Accessed December 14, 2020. Available at URL address: <http://www.nice.org.uk/guidance/CG61>
79. New Zealand Guidelines Group. Identification of Common Mental Disorders and Management of Depression in Primary Care. An Evidence-based Best Practice Guideline. Published by New Zealand Guidelines Group; Wellington: 2008. Accessed December 14, 2020. Available at URL address: http://www.nzgg.org.nz/search?tag_id=28
80. O'Mathúna DP, Ashford RL. Therapeutic touch for healing acute wounds. *Cochrane Database of Systematic Reviews* 2014, Issue 7. Art. No.: CD002766. DOI: 10.1002/14651858.CD002766.pub3.
81. Panebianco M, Sridharan K, Ramaratnam S. Yoga for epilepsy. *Cochrane Database of Systematic Reviews* 2015, Issue 5. Art. No.: CD001524. DOI: 10.1002/14651858.CD001524.pub2.

Complimentary and Alternative Medicine (CAM), continued

83. Passalacqua G. ARIA update: I – Systematic review of complementary and alternative medicine for rhinitis and asthma. *J Allergy Clin Immunol*. 2006 May 1; 117(5): 1054-62.
84. Patyar S, Joshi R, Byrav DS, Prakash A, Medhi B, Das BK. Bacteria in cancer therapy: a novel experimental strategy. *J Biomed Sci*. 2010 Mar 23;17(1):21.
85. Pereira LM, Obara K, Dias JM, Menacho MO, Guariglia DA, Schiavoni D, Pereira HM, Cardoso JR. Comparing the Pilates method with no exercise or lumbar stabilization for pain and functionality in patients with chronic low back pain: systematic review and meta-analysis. *Clin Rehabil*. 2012 Jan;26(1):10-20.
86. Posadzki P, Alotaibi A, Ernst E. Adverse effects of homeopathy: a systematic review of published case reports and case series. *Int J Clin Pract*. 2012 Dec;66(12):1178-88. doi: 10.1111/ijcp.12026.
87. Posadzki P, Lewandowski W, Terry R, Ernst E, Stearns A. Guided imagery for non-musculoskeletal pain: a systematic review of randomized clinical trials. *J Pain Symptom Manage*. 2012 Jul;44(1):95-104. doi: 10.1016/j.jpainsymman.2011.07.014.
88. Rada G, Capurro D, Pantoja T, Corbalán J, Moreno G, Letelier LM, Vera C. Non-hormonal interventions for hot flushes in women with a history of breast cancer. *Cochrane Database of Systematic Reviews* 2010, Issue 9. Art. No.: CD004923. DOI: 10.1002/14651858.CD004923.pub2.
89. Rosche B1, Wemecke KD, Ohlraun S, Dörr JM, Paul F. *Trichuris suis ova* in relapsing-remitting multiple sclerosis and clinically isolated syndrome (TRiOMS): study protocol for a randomized controlled trial. *Trials*. 2013 Apr 25; 14:112.
90. Rothgangel AS, Braun SM, Beurskens AJ, Seitz RJ, Wade DT. The clinical aspects of mirror therapy in rehabilitation: a systematic review of the literature. *Int J Rehabil Res*. 2011 Mar;34(1):1-13.
91. Royal Australian and New Zealand College of Obstetrician and Gynaecologists. Vitamin and mineral supplementation in pregnancy. Nov 2014. Accessed December 14, 2020. Available at URL address: <http://www.ranzcog.edu.au/college-statements-guidelines.html>
92. Sandborn WJ1, Elliott DE, Weinstock J, Summers RW, Landry-Wheeler A, Silver N, Harnett MD, Hanauer SB. Randomised clinical trial: the safety and tolerability of *Trichuris suis ova* in patients with Crohn's disease. *Aliment Pharmacol Ther*. 2013 Aug;38(3):255-63.
93. Sawitzke AD, Shi H, Finco MF, Dunlop DD, Harris CL, Singer NG, Bradley JD, Silver D, Jackson CG, Lane NE, Oddis CV, Wolfe F, Lisse J, Furst DE, Bingham CO, Reda DJ, Moskowitz RW, Williams HJ, Clegg DO. Clinical efficacy and safety of glucosamine, chondroitin sulphate, their combination, celecoxib or placebo taken to treat osteoarthritis of the knee: 2-year results from GAIT. *Ann Rheum Dis*. 2010 Aug;69(8):1459-64.
94. Selles RW, Schreuders TA, Stam HJ. Mirror therapy in patients with causalgia (complex regional pain syndrome type II) following peripheral nerve injury: two cases. *J Rehabil Med*. 2008 Apr;40(4):312-4.
95. Sephton SE, Salmon P, Weissbecker I, Ulmer C, Floyd A, Hoover K, Studts JL. Mindfulness meditation alleviates depressive symptoms in women with fibromyalgia: results of a randomized clinical trial. *Arthritis Rheum*. 2007 Feb 15;57(1):77-85.
96. Shen YH, Nahas R. Complementary and alternative medicine for treatment of irritable bowel syndrome. *Can Fam Physician*. 2009 Feb;55(2):143-8.
97. Siniscalco D1, Antonucci N. Possible use of *Trichuris suis ova* in autism spectrum disorders therapy. *Med Hypotheses*. 2013 Jul;81(1):1-4.
98. Skillgate E, Bohman T, Holm LW, Vingård E, Alfredsson L. The long-term effects of naprapathic manual therapy on back and neck pain - results from a pragmatic randomized controlled trial. *BMC Musculoskelet Disord*. 2010 Feb 5; 11:26.
99. Smith CA, Collins CT, Crowther CA. Aromatherapy for pain management in labour. *Cochrane Database of Systematic Reviews*. 2011, Issue 7. Art. No.: CD009215. DOI: 10.1002/14651858.CD009215.
100. Smith CA, Collins CT, Cyna AM, Crowther CA. Complementary and alternative therapies for pain management in labour. *Cochrane Database of Systematic Reviews* 2006, Issue 4. Art. No.: CD003521. DOI: 10.1002/14651858.CD003521.pub2.
101. Smith CA, Levett KM, Collins CT, Jones L. Massage, reflexology and other manual methods for pain management in labour. *Cochrane Database of Systematic Reviews* 2012, Issue 2. Art. No.: CD009290. DOI: 10.1002/14651858.CD009290.pub2.
102. Smith N, Weymann A, Tausk FA, Gelfand JM. Complementary and alternative medicine for psoriasis: a qualitative review of the clinical trial literature. *J Am Acad Dermatol*. 2009 Nov;61(5):841-56.
103. Summers RW. Novel and future medical management of inflammatory bowel disease. *Surg Clin North Am*. 2007 Jun;87(3):727-41.
104. Summers RW, Elliott DE, Urban JF Jr, Thompson RA, Weinstock JV. *Trichuris suis* therapy for active ulcerative colitis: a randomized controlled trial. *Gastroenterology*. 2005 Apr;128(4):825-32.
105. Szczerko O, Cooley K, Mills EJ, Zhou Q, Pemi D, Seely D. Naturopathic treatment of rotator cuff tendinitis among Canadian postal workers: a randomized controlled trial. *Arthritis Rheum*. 2009 Aug 15;61(8):1037-45.
106. Teut M, Lüdtkke R, Schnabel K, Willich SN, Witt CM. Homeopathic treatment of elderly patients—a prospective observational study with follow-up over a two year period. *BMC Geriatr*. 2010 Feb 22; 10:10.
107. Thachil AF, Mohan R, Bhugra D. The evidence base of complementary and alternative therapies in depression. *J Affect Disord*. 2007 Jan;97(1-3):23-35.
108. Thieme H, Mehrholz J, Pohl M, Behrens J, Dohle C. Mirror therapy for improving motor function after stroke. *Cochrane Database of Systematic Reviews* 2012, Issue 3. Art. No.: CD008449. DOI: 10.1002/14651858.CD008449.pub2.
109. U.S. Food and Drug Administration (FDA). Draft guidance. Complementary and alternative medicine products and their regulation by the Food and Drug Administration. Mar 2, 2007. Accessed Jun 4, 2015. Available at URL address: <http://www.fda.gov/RegulatoryInformation/Guidances/ucm144657.htm>
110. U.S. Food and Drug Administration (FDA). Labeling & nutrition. Food labeling and nutritional overview. Feb 27, 2014. Accessed Jun 4, 2015. Available at URL address: <http://www.fda.gov/food/ingredientspackaginglabeling/labelingnutrition/default.htm>
111. Vancampfort D, Vansteelandt K, Scheewe T, Probst M, Knapen J, De Herdt A, De Hert M. Yoga in schizophrenia: a systematic review of randomised controlled trials. *Acta Psychiatr Scand*. 2012 Jul;126(1):12-20.
112. VanderVaart S, Gijzen VM, de Wildt SN, Koren G. A systematic review of the therapeutic effects of Reiki. *J Altern Complement Med*. 2009 Nov;15(11):1157-69.
113. Verma S, Thuluvath PJ. Complementary and alternative medicine in hepatology: review of the evidence of efficacy. *Clin Gastroenterol Hepatol*. 2007 Apr;5(4):408-16.
114. Wang C, Bannuru R, Ramel J, Kupelnick B, Scott T, Schmid CH. Tai Chi on psychological well-being: systematic review and meta-analysis. *BMC Complement Altern Med*. 2010 May 21; 10:23.
115. Wang CW, Ng SM, Ho RT, Ziea ET, Wong VC, Chan CL. The effect of qigong exercise on immunity and infections: a systematic review of controlled trials. *Am J Chin Med*. 2012;40(6):1143-56. doi: 10.1142/S0192415X1250084X.
116. Webb AN, Kukuruzovic RH, Catto-Smith AG, Sawyer SM. Hypnotherapy for treatment of irritable bowel syndrome. *Cochrane Database of Systematic Reviews* 2007, Issue 4. Art. No.: CD005110. DOI: 10.1002/14651858.CD005110.pub2.

General Medicine Policies, Continued

Complimentary and Alternative Medicine (CAM), continued

117. Wilkens P, Scheel IB, Grundnes O, Hellum C, Storheim K. Effect of glucosamine on pain-related disability in patients with chronic low back pain and degenerative lumbar osteoarthritis: a randomized controlled trial. *JAMA*. 2010 Jul 7;304(1):45-52.
118. Wołowiec P, Michalak I, Chojnacka K, Mikulewicz M. Hair analysis in health assessment. *Clin Chim Acta*. 2013 Apr 18;419:139-71. doi: 10.1016/j.cca.2013.02.001.
119. Wood MJ, Molassiotis A, Payne S. What research evidence is there for the use of art therapy in the management of symptoms in adults with cancer? A systematic review. *Psychooncology*. 2011 Feb;20(2):135-45. doi: 10.1002/pon.1722.
120. World Health Organization (WHO). WHO traditional medicine strategy: 2014-2023 Accessed December 14, 2020. Available at URL address: http://www.who.int/medicines/publications/traditional/trm_strategy14_23/en/
121. World Health Organization (WHO). Legal status of traditional medicine and complementary/alternative medicine: A worldwide review. Accessed Jun 4, 2015. Available at URL address: <http://apps.who.int/medicinedocs/en/d/Jh2943e/>
122. Yeh GY, Wang C, Wayne PM, Phillips R. Tai chi exercise for patients with cardiovascular conditions and risk factors: A SYSTEMATIC REVIEW. *J Cardiopulm Rehabil Prev*. 2009 May-Jun;29(3):152-60.
123. Yun GJ, Chun MH, Park JY, Kim BR. The synergic effects of mirror therapy and neuromuscular electrical stimulation for hand function in stroke patients. *Ann Rehabil Med*. 2011 Jun;35(3):316-21.
124. Zeng X, Liu M, Yang Y, Li Y, Asplund K. Ginkgo biloba for acute ischaemic stroke. *Cochrane Database of Systematic Reviews* 2005, Issue 4. Art. No.: CD003691. DOI: 10.1002/14651858.CD003691.pub2
125. Zhang M, Liu X, Li J, He L, Tripathy D. Chinese medicinal herbs to treat the side-effects of chemotherapy in breast cancer patients. *Cochrane Database of Systematic Reviews* 2007, Issue 2. Art. No.: CD004921. DOI: 10.1002/14651858.CD004921.pub2.

Disclaimer

This document is for informational purposes only and should not be relied on in the diagnosis and care of individual patients. Medical and Coding/Reimbursement policies do not constitute medical advice, plan preauthorization, certification, an explanation of benefits, or a contract. Members should consult with appropriate healthcare providers to obtain needed medical advice, care, and treatment. Benefits and eligibility are determined before medical guidelines and payment guidelines are applied. Benefits are determined by the member's individual benefit plan that is in effect at the time services are rendered.

The codes for treatments and procedures applicable to this policy are included for informational purposes. Inclusion or exclusion of a procedure, diagnosis or device code(s) does not constitute or imply member coverage or provider reimbursement policy. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.

SelectHealth® makes no representations and accepts no liability with respect to the content of any external information cited or relied upon in this policy. SelectHealth updates its Coverage Policies regularly, and reserves the right to amend these policies without notice to healthcare providers or SelectHealth members.

Members may contact Customer Service at the phone number listed on their member identification card to discuss their benefits more specifically. Providers with questions about this Coverage Policy may call SelectHealth Provider Relations at (801) 442-3692.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from SelectHealth.

"Intermountain Healthcare" and its accompanying logo, the marks of "SelectHealth" and its accompanying marks are protected and registered trademarks of the provider of this Service and or Intermountain Health Care, Inc., IHC Health Services, Inc., and SelectHealth, Inc. Also, the content of this Service is proprietary and is protected by copyright. You may access the copyrighted content of this Service only for purposes set forth in these Conditions of Use.

© CPT Only – American Medical Association



LONG-TERM ACUTE CARE (LTAC)

Policy # 583

Implementation Date: 7/20/16

Review Dates: 6/15/17, 7/16/18, 6/17/19, 6/14/20, 6/16/21

Revision Dates:

Related Medical Policies:

[#443 Acute Inpatient Rehabilitation](#)

Disclaimer:

1. Policies are subject to change without notice.
2. Policies outline coverage determinations for SelectHealth Commercial, SelectHealth Advantage (Medicare/CMS), and SelectHealth Community Care (Medicaid/CHIP) plans. Refer to the "Policy" section for more information.

Description

Long-term acute care hospitals [LTACH] (also called transitional care hospitals) are licensed as acute care hospitals with additional Medicare certification that supports a length of stay measured in weeks (more than 25 days on average for Medicare patients) as compared to the typical five-day stay for patients in traditional hospitals. LTACH are somewhat unique in their focus on care for critically ill patients, who require specialized, aggressive, goal-directed care over an extended recovery period. LTAC hospitals provide long-term acute care (LTAC) to complex medically complex patients who require an extended stay in a hospital setting. Typical patients have multiple co-morbidities, multi organ system failure, and significant loss of independence, most following a traditional hospital stay.

These hospitals are exempt from the APR DRG methodology and are reimbursed a hospital specific rate paid per day of covered inpatient care. An example of a service provided by a long-term stay hospital is ventilator care. The term "long-term stay hospital" does not include a psychiatric, rehabilitation, or children's hospital. These facilities may be free-standing or part of a general acute care facility.

Commercial Plan Policy/CHIP (Children's Health Insurance Program)

Application of coverage criteria is dependent upon an individual's benefit coverage at the time of the request.

SelectHealth **covers long-term acute care (LTAC)** as medically necessary **when specific criteria are met.**

Coverage Criteria (Must meet either 1 or 2):

1. **Complex Medical Needs with Significant Functional Impairment(s): (must have all)**
 - a. Member is admitted directly from an inpatient hospital, but the discharge was not from a psychiatric or rehabilitation facility.
 - b. Documentation of an established diagnosis or condition for which ongoing acute hospital care is needed.
 - c. Documentation that indicates the member will benefit from and improve with the LTAC program available at the chosen facility.
 - d. Documentation indicates an expectation that the member will require long-term acute care for a length of stay of 30 or more days.
 - e. Requires daily medical practitioner assessment or intervention.

General Medicine Policies, Continued

Long-Term Acute Care (LTAC), continued

- f. Care at LTACH is appropriate for condition as indicated by InterQual criteria.
2. **Ventilator Management and Weaning - Admission to LTACH may be medically necessary when ALL the following are present:**
- a. Requires daily medical practitioner assessment or intervention
 - b. InterQual Criteria for ventilator management and weaning are met

Discharge for NON-ventilator patients from the LTAC facility is appropriate when (ALL must be met):

- a. The member is hemodynamically stable without daily medication adjustments
- b. The member no longer requires multiple intravenous drug therapy
- c. The member no longer requires cardiac monitoring
- d. The member has a stable hemoglobin and hematocrit without transfusion and stable electrolytes without daily parenteral adjustments
- e. The member is stable on current nutritional support (whether it is parenteral, oral, or percutaneous G/J tube)
- f. The member no longer requires hemodialysis or is stable for transport to and from hemodialysis
- g. The member is able to participate in, but is not receiving, at least 3 hours of therapy daily
- h. All care including wound care can be managed at a lower level of care

Discharge for Ventilator Patients from the LTAC facility is appropriate when:

- a. The member is hemodynamically stable without daily medication adjustment
- b. The member is stable off the ventilator or is stable on the ventilator and considered not able to be weaned
- c. Is clear of infection or is stable on antibiotic regimen
- d. All care can be managed at a lower level of care

SelectHealth Advantage (Medicare/CMS)

Coverage is determined by the Centers for Medicare and Medicaid Services (CMS); if a coverage determination has not been adopted by CMS, and InterQual criteria are not available, the SelectHealth Commercial policy applies. For the most up-to-date Medicare policies and coverage, please visit their search website <http://www.cms.gov/medicare-coverage-database/overview-and-quick-search.aspx?from2=search1.asp&> or the [manual website](#)

SelectHealth Community Care (Medicaid)

SelectHealth does NOT cover LTAC for managed Medicaid members as typical stay is greater than or equal to 30 days and member enrollment is automatically transferred to fee for service Medicaid.

General Medicine Policies, Continued

Long-Term Acute Care (LTAC), continued

Billing/Coding Information

CPT CODES

No specific codes identified

HCPCS CODES

No specific codes identified

Key References

1. Centers for Medicaid and Medicare Services. CMS Long-Term Care Hospital PPS Overview at http://www.cms.hhs.gov/LongTermCareHospitalPPS/03_ltch_train.asp#TopOfPage.
2. CMS Long-Term Care Hospital PPS Overview at http://www.cms.hhs.gov/LongTermCareHospitalPPS/03_ltch_train.asp#TopOfPage.
3. Harvard Pilgrim HealthCare "Medical Review Criteria Inpatient Rehabilitation/LongTerm Acute Care," effective 4/12/2012 UnitedHealthcare Policy H-SHO-006, "Hospital Services (Inpatient and Outpatient)," revised 9/7/2010.
4. Illinois Health and Family Services. Handbook for Providers of Hospital Services, Chapter H-200. Version date. September 2014. Retrieved from: <http://www2.illinois.gov/hfs/SiteCollectionDocuments/h200.pdf>
5. Iowa Department of Human Services. Physician Services Provider Manual. Version date, July 1, 2014. Retrieved from: <https://dhs.iowa.gov/sites/default/files/Phys.pdf>
6. McKesson. (2011). Alternate level of care guidelines: long-term acute care. Retrieved April 3, 2012 from CareEnhance Review Manager database.
7. Michigan Department of Community Health. Medicaid Provider Manual. Version date, October 1, 2014. Retrieved from: <http://www.mdch.state.mi.us/dch-medicaid/manuals/MedicaidProviderManual.pdf>
8. Munoz-Price, L. S. (2009). Long term acute care hospital. *Clinical Infectious Diseases*, 49 (3), 438-43.
9. Pate, N.P., & Malagoni, M.A. (2009). Antimicrobial agents for surgical infections. *Surgical Clinics of North America*, 89(3), 611-626. DOI:10.1016/j.suc.2009.03.009
10. Pediatric Section Task Force on Admission and Discharge Criteria, Society of Critical Care Medicine in conjunction with the American College of Critical Care Medicine and the Committee on Hospital Care of the American Academy of Pediatrics. (2007). Guidelines for developing admission and discharge policies for the pediatric intensive care unit. *Critical Care Medicine*, 27(4), 843-845.
11. WellCare Policy HS-162, "Long Term Acute Care Hospital: Criteria for Admission," revised 6/7/12.

Disclaimer

This document is for informational purposes only and should not be relied on in the diagnosis and care of individual patients. Medical and Coding/Reimbursement policies do not constitute medical advice, plan preauthorization, certification, an explanation of benefits, or a contract. Members should consult with appropriate healthcare providers to obtain needed medical advice, care, and treatment. Benefits and eligibility are determined before medical guidelines and payment guidelines are applied. Benefits are determined by the member's individual benefit plan that is in effect at the time services are rendered.

The codes for treatments and procedures applicable to this policy are included for informational purposes. Inclusion or exclusion of a procedure, diagnosis or device code(s) does not constitute or imply member coverage or provider reimbursement policy. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.

SelectHealth® makes no representations and accepts no liability with respect to the content of any external information cited or relied upon in this policy. SelectHealth updates its Coverage Policies regularly, and reserves the right to amend these policies without notice to healthcare providers or SelectHealth members.

Members may contact Customer Service at the phone number listed on their member identification card to discuss their benefits more specifically. Providers with questions about this Coverage Policy may call SelectHealth Provider Relations at (801) 442-3692.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from SelectHealth.

"Intermountain Healthcare" and its accompanying logo, the marks of "SelectHealth" and its accompanying marks are protected and registered trademarks of the provider of this Service and or Intermountain Health Care, Inc., IHC Health Services, Inc., and SelectHealth, Inc. Also, the content of this Service is proprietary and is protected by copyright. You may access the copyrighted content of this Service only for purposes set forth in these Conditions of Use.

© CPT Only – American Medical Association



MEDICAL TRANSPORTATION

Policy # 344

Implementation Date: 7/19/98

Review Dates: 1/04/00, 2/27/01, 3/29/02, 5/17/07, 4/24/08, 4/23/09, 4/21/11, 4/12/12, 8/15/13, 6/19/14, 6/11/15, 6/16/16, 6/15/17, 6/20/19, 6/15/20, 6/17/21, 6/30/22

Revision Dates: 2/18/10, 10/7/21, 9/7/22

Disclaimer:

1. Policies are subject to change without notice.
2. Policies outline coverage determinations for SelectHealth Commercial, SelectHealth Advantage (Medicare/CMS), and SelectHealth Community Care (Medicaid/CHIP) plans. Refer to the “Policy” section for more information.

Description

SelectHealth members may require medical transportation when other means of transportation would endanger the patient's health, or when there are no other feasible methods of transportation available. In these instances, the involved providers or members may request coverage of medical transportation in conjunction with applicable medical benefits.

Commercial Plan Policy/CHIP (Children's Health Insurance Program)

Application of coverage criteria is dependent upon an individual's benefit coverage at the time of the request.

SelectHealth **provides coverage of medical transport when the following criteria are met:**

Non-emergent Medical Transportation:

Inpatient facility-to-facility medical transport is covered when:

1. The member is currently admitted at a non-contracted facility, or at a contracted facility that is unable to meet the member's medical needs, or at a contracted facility that is greater than 50 miles from member's home and member will require post-acute follow-up; **and**
2. The member cannot be safely transported through non-medical transportation (such as a car or commercial airline); **and**
3. The method of transportation requested is the minimum necessary or most efficient to safely meet the member's medical needs; **and**
4. Discharge is not imminent from current facility.

Examples of non-emergency medical transport include, but are not limited to:

- Inter-facility transfers
- Transfers between hospitals, skilled nursing facilities, or rehabilitation centers
- Transportation for inpatient tests not available at the admitting facility
- Round trip transportation from one facility to another (including a physician's office) to obtain medically necessary diagnostic or therapeutic services

General Medicine Policies, Continued

Medical Transportation, continued

Emergent or Urgent Medical Transportation:

Emergency medical transport does not require preauthorization. Refer to the 'Ambulance & Paramedics' Benefit Clarification for more details.

Emergent Air Transportation

The member's medical condition requires immediate and rapid transport, which cannot be sufficiently provided by ground ambulance, or the member's location is inaccessible by ground ambulance; wherein, in either situation, this would pose a threat to the member's survival or endanger their health.

Limitations/Exclusions:

- Inter-facility transfers due to member preference or convenience.
- Outpatient transportation for non-emergent physician office visits or treatments.
- Death-related transportation following the pronouncement of death by a person legally authorized to make this determination (MD, State Medical Examiner) is excluded.
- Transportation to the coroner's office or mortuary.

SelectHealth Advantage (Medicare/CMS)

Coverage is determined by the Centers for Medicare and Medicaid Services (CMS); if a coverage determination has not been adopted by CMS, and InterQual criteria are not available, the SelectHealth Commercial policy applies. For the most up-to-date Medicare policies and coverage, please visit their search website <http://www.cms.gov/medicare-coverage-database/overview-and-quick-search.aspx?from2=search1.asp&> or [the manual website](#)

SelectHealth Community Care (Medicaid)

Coverage is determined by the State of Utah Medicaid program; if Utah State Medicaid has no published coverage position and InterQual criteria are not available, the SelectHealth Commercial criteria will apply. For the most up-to-date Medicaid policies and coverage, please visit their website <http://health.utah.gov/medicaid/manuals/directory.php> or the [Utah Medicaid code Look-Up tool](#)

Billing/Coding Information

Covered: For the indications outlined above

CPT CODES

No specific codes identified

HCPCS CODES

A0425	Ground mileage, per statute mile
A0426	Ambulance service, advanced life support, non-emergency transport, level 1 (ALS 1)
A0427	Ambulance service, advanced life support, emergency transport, level 1 (ALS1-emergency)
A0428	Ambulance service, basic life support, non-emergency transport, (BLS)
A0429	Ambulance service, basic life support, emergency transport (BLS-emergency)
A0430	Ambulance service, conventional air services, transport, one way (fixed wing)



WOUND CARE AND PHYSICAL THERAPY

Policy # 469

Implementation Date: 12/20/10

Review Dates: 12/15/11, 7/18/13, 8/28/14, 8/20/15, 8/25/16, 8/17/17, 7/25/18, 6/14/19, 6/4/20, 6/17/21

Revision Dates: 6/20/19

Disclaimer:

1. Policies are subject to change without notice.
2. Policies outline coverage determinations for SelectHealth Commercial, SelectHealth Advantage (Medicare/CMS), and SelectHealth Community Care (Medicaid/CHIP) plans. Refer to the "Policy" section for more information.

Description

The intent of the physical therapy limit is to impose a cap on traditional physical therapy services post-injury due to the significant variability that occurs from provider to provider and patient to patient for the same condition. It is also intended to promote the responsible use of this limited benefit by patients and providers. Additionally, actuarial analysis has shown that most individuals will not use their maximum number of visits in a plan year, and thus, this limitation should serve nearly every member's need.

Wound care may present a circumstance in which application of the benefit limitation may not serve the plan or the member. Wound care is often performed by a physical therapist in an outpatient setting. As such, these services are billed using physical therapy CPT codes consistent with the services provided and the specialty billing the service. In some cases, especially chronic diabetic or large wounds, wound care/therapy may occur over many months, and thus, patients may meet their plan limitations for physical therapy quite rapidly. Thus, a process needs to be in place which allows for consistent decision-making in allowing for exceptions to the coverage limitation based upon the plan's discretionary authority granted in the certificate of coverage in situations in which it is financially prudent to offer the exception.

Commercial Plan Policy/CHIP (Children's Health Insurance Program)

Application of coverage criteria is dependent upon an individual's benefit coverage at the time of the request.

SelectHealth **covers extended physical therapy benefits in the management of chronic wounds in *limited circumstances*.**

Criteria for coverage:

1. Documentation includes medical necessity for utilizing a physical therapist to provide wound care rather than a trained wound care nurse or technician.
2. Documentation indicates that all services being provided by the physical therapist are directly related to wound care only; if the member is receiving wound care therapy in addition to other non-wound related physical therapy, only the wound care portions of the therapy will be applied toward the benefit limit exception.
3. Documentation must demonstrate progress toward healing the wound. This is defined as a reduction in the wound size/volume of at least 10% for every month in which wound care is performed.
4. Services being provided are a covered benefit and not investigational/experimental by the plan's definition.
5. The requesting provider must submit a plan of therapy with identified outcomes and goals.

General Medicine Policies, Continued

Wound Care and Physical Therapy, continued

6. Any exception which is approved is for a maximum of 90 days. Further exceptions require reassessment by clinical reviewers.
7. Member is being case managed by a SelectHealth RN Care Manager.

SelectHealth Advantage (Medicare/CMS)

Coverage is determined by the Centers for Medicare and Medicaid Services (CMS); if a coverage determination has not been adopted by CMS, and InterQual criteria are not available, the SelectHealth Commercial policy applies. For the most up-to-date Medicare policies and coverage, please visit their search website <http://www.cms.gov/medicare-coverage-database/overview-and-quick-search.aspx?from2=search1.asp&> or [the manual website](#)

SelectHealth Community Care (Medicaid)

Coverage is determined by the State of Utah Medicaid program; if Utah State Medicaid has no published coverage position and InterQual criteria are not available, the SelectHealth Commercial criteria will apply. For the most up-to-date Medicaid policies and coverage, please visit their website <http://health.utah.gov/medicaid/manuals/directory.php> or the [Utah Medicaid code Look-Up tool](#)

Summary of Medical Information

Many things can prevent a wound from healing, such as diseases (e.g., diabetes, poor blood flow, or other vascular problems) medications, infections, and/or poor nutrition. A wound that doesn't heal places a person at risk for infection and more serious complications. The wound can cause discomfort, pain, disfigurement, and can also place limits on a person's activities and quality of life.

Wound clinics provide treatment for chronic or non-healing wounds. The clinic provides a comprehensive, individualized plan, indicating the wound problem and goal of treatment. The types of wounds treated include non-healing surgical wounds, pressure ulcers, arterial ulcers, diabetic related wounds and ulcers, venous insufficiency, and burns.

Wound clinics use a multi-disciplinary approach to wound care by bringing together a team of specially trained doctors, nurses, and other clinicians experienced in all aspects of wound management. They can identify compounding factors such as malnutrition, metabolic disorders, and structural abnormalities that can retard normal healing and predispose the patient for recurrence. An aggressive approach to therapy can aid in early intervention, improved healing rates, reduced amputations, and reduced disability.

Wound care management improves the patient's mobility, quality of life, and overall well-being. Wounds are treated based on cause and severity using a variety of wound care products to support the healing process. In patients suffering from chronic wounds, wound care management has become critical in reducing the risk to the body, which can, in some cases, include loss of life.

The treatment of chronic wounds, such as pressure ulcers, requires prolonged care. During the prolonged treatment process, the patient remains at risk of developing new pressure ulcers in other places, so the treatment is based on the staging of the pressure ulcer. Debridement, topical wound care, treatment of infection, control of chronic wound contamination, the positioning of the patient, and the use of support surfaces all play an important role in the treatment of pressure ulcers.

Venous ulcers are treated using compression therapy, debridement of necrotic tissue, and by providing a moist wound environment. Some venous ulcers are treated using the application of bioengineered skin or split-thickness skin grafting. The treatment of diabetic foot ulcers requires control of blood glucose levels, appropriate footwear, saline dressings to provide a moist wound environment, debridement and evaluation, and the correction of peripheral arterial insufficiency. Negative pressure wound therapy, heterogenic dressings and/or grafts, and the application of cellular tissue products may also be employed to assist in wound healing.

Technological advances such as improvements in synthetic dressing materials and newer technologies, such as recombinant growth factors, endovascular arterial and venous repair techniques, epidermal grafting, bi-layered human dermal substitutes, cellular tissue products, and xenogeneic tissue scaffolds

General Medicine Policies, Continued

Wound Care and Physical Therapy, continued

are expected to have a positive impact on wound care management. The introduction of new and innovative technologies into this area is helping to meet patient demands. Additionally, techniques such as electrical stimulation, electromagnetic therapy, nanotechnology, therapeutic ultrasound, and the use of silver and combination dressings offer the potential for further improving the way wound care is managed.

Billing/Coding Information

Covered: *For the conditions outlined above*

CPT CODES

- 97161** Physical therapy evaluation: low complexity, requiring these components: A history with no personal factors and/or comorbidities that impact the plan of care; An examination of body system(s) using standardized tests and measures addressing 1-2 elements from any of the following: body structures and functions, activity limitations, and/or participation restrictions; A clinical presentation with stable and/or uncomplicated characteristics; and Clinical decision making of low complexity using standardized patient assessment instrument and/or measurable assessment of functional outcome. Typically, 20 minutes are spent face-to-face with the patient and/or family.
- 97162** Physical therapy evaluation: moderate complexity, requiring these components: A history of present problem with 1-2 personal factors and/or comorbidities that impact the plan of care; An examination of body systems using standardized tests and measures in addressing a total of 3 or more elements from any of the following: body structures and functions, activity limitations, and/or participation restrictions; An evolving clinical presentation with changing characteristics; and Clinical decision making of moderate complexity using standardized patient assessment instrument and/or measurable assessment of functional outcome. Typically, 30 minutes are spent face-to-face with the patient and/or family.
- 97163** Physical therapy evaluation: high complexity, requiring these components: A history of present problem with 3 or more personal factors and/or comorbidities that impact the plan of care; An examination of body systems using standardized tests and measures addressing a total of 4 or more elements from any of the following: body structures and functions, activity limitations, and/or participation restrictions; A clinical presentation with unstable and unpredictable characteristics; and Clinical decision making of high complexity using standardized patient assessment instrument and/or measurable assessment of functional outcome. Typically, 45 minutes are spent face-to-face with the patient and/or family.
- 97164** Re-evaluation of physical therapy established plan of care, requiring these components: An examination including a review of history and use of standardized tests and measures is required; and revised plan of care using a standardized patient assessment instrument and/or measurable assessment of functional outcome typically, 20 minutes are spent face-to-face with the patient and/or family.
- 97022** Application of a modality to one or more areas; whirlpool
- 97024** ; diathermy (e.g., microwave)
- 97026** ; infrared
- 97028** ; ultraviolet
- 97032** ; electrical stimulation (manual), each 15 minutes
- 97033** ; iontophoresis, each 15 minutes
- 97035** ; ultrasound, each 15 minutes
- 97036** ; Hubbard tank, each 15 minutes
- 97597** Debridement (eg, high pressure waterjet with/without suction, sharp selective debridement with scissors, scalpel and forceps), open wound, (eg, fibrin, devitalized epidermis and/or dermis, exudate, debris, biofilm), including topical application(s), wound assessment, use of a whirlpool, when performed and instruction(s) for ongoing care, per session, total wound(s) surface area; first 20 sq cm or less

General Medicine Policies, Continued

Wound Care and Physical Therapy, continued

- 97598** ; each additional 20 sq cm, or part thereof (List separately in addition to code for primary procedure)
- 97602** Removal of devitalized tissue from wound(s), non-selective debridement, without anesthesia (e.g., wet-to-moist dressings, enzymatic, abrasion), including topical application(s), wound assessment, and instruction(s) for ongoing care, per session
- 97605** Negative pressure wound therapy (e.g., vacuum assisted drainage collection), including topical application(s), wound assessment, and instruction(s) for ongoing care, per session; total wound(s) surface area less than or equal to 50 square centimeters
- 97606** ; total wound(s) surface area greater than 50 square centimeters

HCPCS CODES

- G0281** Electrical stimulation, (unattended), to one or more areas, for chronic Stage III and Stage IV pressure ulcers, arterial ulcers, diabetic ulcers, and venous stasis ulcers not demonstrating measurable signs of healing after 30 days of conventional care, as part of a therapy plan of care
- G0282** Electrical stimulation, (unattended), to one or more areas, for wound care other than described in G0281
- G0295** Electromagnetic therapy, to one or more areas, for wound care other than described in G0329 or for other uses
- G0329** Electromagnetic therapy, to one or more areas for chronic Stage III and Stage IV pressure ulcers, arterial ulcers, diabetic ulcers and venous stasis ulcers not demonstrating measurable signs of healing after 30 days of conventional care as part of a therapy plan of care

Key References

1. Agency for Health Care Research and Quality (AHRQ). Treatment of pressure ulcers. Clinical guide, No.15. Rockville, MD: U.S. Department of Health and Human Services, Public Health Service, Agency for Health Care Policy and Research (AHCPR), publication no. 95-0652, 1994.
2. *American Physical Therapy Association. Guide to Physical Therapist Practice. 2nd ed.* Phys Ther, 2001; 81:9-744.
3. Stillman RM, Daley BJ, Talavera et al. Wound Care. Updated April 6, 2010. Available: <http://emedicine.medscape.com/article/194018-treatment>. Date Accessed: November 16, 2010.
4. Vittal, B. (2010). A Growing Market: Wound Care Management. Hospital Management. Updated: October 6, 2010. Available: <http://www.hospitalmanagement.net/features/feature97878/>. Date Accessed: November 20, 2010.

Disclaimer

This document is for informational purposes only and should not be relied on in the diagnosis and care of individual patients. Medical and Coding/Reimbursement policies do not constitute medical advice, plan preauthorization, certification, an explanation of benefits, or a contract. Members should consult with appropriate healthcare providers to obtain needed medical advice, care, and treatment. Benefits and eligibility are determined before medical guidelines and payment guidelines are applied. Benefits are determined by the member's individual benefit plan that is in effect at the time services are rendered.

The codes for treatments and procedures applicable to this policy are included for informational purposes. Inclusion or exclusion of a procedure, diagnosis or device code(s) does not constitute or imply member coverage or provider reimbursement policy. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.

SelectHealth® makes no representations and accepts no liability with respect to the content of any external information cited or relied upon in this policy. SelectHealth updates its Coverage Policies regularly, and reserves the right to amend these policies without notice to healthcare providers or SelectHealth members.

Members may contact Customer Service at the phone number listed on their member identification card to discuss their benefits more specifically. Providers with questions about this Coverage Policy may call SelectHealth Provider Relations at (801) 442-3692.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from SelectHealth.

"Intermountain Healthcare" and its accompanying logo, the marks of "SelectHealth" and its accompanying marks are protected and registered trademarks of the provider of this Service and or Intermountain Health Care, Inc., IHC Health Services, Inc., and SelectHealth, Inc. Also, the content of this Service is proprietary and is protected by copyright. You may access the copyrighted content of this Service only for purposes set forth in these Conditions of Use.

© CPT Only – American Medical Association